

300
17
15083
1

JUN 28 1985

PureGro Company
1276 Halyard Drive
West Sacramento, CA 95691

Attention: Christopher L. Dozier

Gentlemen:


Subject: PureGro Liquid Copper Fungicide
EPA Registration No. 1202-301
Your Letter Dated December 27, 1984

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted.

1. Change the heading "Environmental Cautions" to "Environmental Hazards"
2. Change "Keep out of lakes, streams, or ponds" to "Do not apply directly to water."
3. Expand the statement of practical treatment by adding statements applicable to dermal, eye, and inhalation exposure or contact.

A stamped copy of the label is enclosed for your records.

Sincerely yours,


Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

RD/PTB; JOB: 88609; Files: RD: 65/27; Kendrick&Co: 898-1270; 5/31/85; Dw1: 6/6/85; DED

COMPLIANCE

SYMBOL	DATE	INITIALS	REMARKS	DATE	INITIALS	REMARKS	DATE	INITIALS	REMARKS

**HAZARDS TO HUMANS
(AND DOMESTIC ANIMALS)
CAUTION**

Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuff.

STATEMENT OF PRACTICAL TREATMENT

In case of ingestion call a physician immediately. Do Not Induce Vomiting.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Keep Away from Heat, Sparks and Open Flames • Use with Adequate Ventilation.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not apply this product to such a extent as to directly or through drift expose workers or other persons. The area being treated must be roped off or restricted persons, and spray should not be directed into treated areas without protective clothing.

Because certain crops may require more frequent reentry intervals for certain crops treated with this product, consult your State Department of Agriculture for further information.

Workers or other persons should be given at least one warning before they are to be treated with this product. When oral warnings are given, warnings should be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

CAUTION: Area to be treated with this product on date of application. Do not enter without appropriate protective clothing until spray has dried. If considerably exposed, consult the Statement of Practical Treatment portion of the pesticide label for first aid and medical attention recommendations.

BEST AVAILABLE COPY

LIQUID COPPER FUNGICIDE

An Emulsifiable Copper Fungicide.

ACTIVE INGREDIENTS:

Copper salts of fatty and rosin acids48.00%

(Copper as metallic 4%)

Copper derived from TC-90®

INERT INGREDIENTS52.00%
100.00%

EPA Reg. No. 1202-301-AA
EPA EST. NO.

CAUTION

KEEP OUT OF REACH OF CHILDREN

PureGro Liquid Copper Fungicide is compatible with Mancozeb, Captan, Prodan, organic phosphates, and zinc. Observe all cautions and limitations concerning all products used in mixtures. Consult state agricultural experiment station or state extension authorities for additional information as to timing and rate of applications.

DISTRIBUTED BY



WEST SACRAMENTO, CALIFORNIA

NET CONTENT

GALS

Do not contaminate water. PEST Wastes resulting from the on site or at an approved CONT

Then offer for recycling and dispose of in a manner approved by local health department. Do not reuse container.

RECOMMENDATIONS

CITRUS (all) — greasy fruit: Use 1.0 to 2.0 gallons coverage ground application. Use 2.0 to 3.0 gallons per acre for good coverage. Apply at first sign of disease. Use 2.0 quarts per acre for aerial application, or 6.0 to 8.0 quarts per acre for aircraft. Apply 1 to 2 weeks before harvest.

WALNUTS — blight: Use 500 gallons per acre water. Apply 500 gallons per acre beginning in early pre-bloom and repeat 10 to 15 percent of the prebloom and immediately before harvest.

BEANS (dry - colored) — blights (Halo and Common): Use 2.0 quarts per acre for ground equipment application per acre. Begin 2 weeks before first appearance. Repeat 4 or 5 times during season.

CANTALOUPE, WATERMELONS — Powdery mildew: Use 2 quarts/100 gallons per acre. Use 2 quarts/100 gallons per acre if disease is expected to appear. Begin 2 weeks before harvest when first noted. Repeat as needed.

BEST AVAILABLE COPY

LOT NO