DIRECTIONS FOR USE

[illegible] Apply [illegible]

BEANS: Mexican Bean Beetle, Leafhoppers, Powdery Mildew — 15 lbs per acre. Lygus Bugs — 20 lbs per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. May be used up to harvest. Sulfur may injure certain varieties of beans.

CORN: Corn Earworm — 15 to 20 lbs per acre. Apply at 3 day intervals starting when silks first appear. Application of this product to the tassel region of corn during the pollen shed period will severely reduce bee populations. Make 4 to 5 applications until silks dry and turn brown. May be applied up to harvest.

COTTON: Early-Season — Thrips, Cotton Fleahopper — 8 to 15 lbs per acre as soon as insects appear. Mid and Late-Season — Boll Weevil, Bollworm, Pink Bollworm, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator — 20 lbs per acre. Apply when insects appear and repeat at 5 to 7 day intervals. Aphid populations will usually be suppressed when this insecticide is used on a regular schedule of treatments.

GRAPES: Grape Leaf Folder — 1st Brood, 10 to 15 lbs [illegible] before first broad larvae emerge from the first rolls. 2nd and 3rd Brood, 10 to 20 lbs per acre. Grape Leafhopper, Powdery Mildew — 10 lbs per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. Thorough coverage is essential. May be applied up to harvest. This application may be expected to control DDT-resistant leafhoppers. Concord and other labrusca types of grapes may be injured by sulfur.

PEACHES, NECTARINES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Cat-facing Insects, Periodical Cicada — 40-60 lbs per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary, repeat every 7 to 10 days. Powdery mildew, [illegible] — 40 to 60 lbs per acre. Apply at [illegible] shuck split, [illegible] 14 days, [illegible] necessary. Do not apply 1 day before harvest. [illegible] 14 days before harvest for nectarines.

PLUMS, PRUNES, CHERRIES [illegible] Peach Twig Borer, Plum Curculio, Red Banded Leaf Roller, Codling Moth, Eastern Tent Caterpillar, Orchard Tent Caterpillar, Western Tussock Moth — 40 to 60 lbs per acre. Apply when insects appear and repeat as necessary. Make full coverage applications. Do not apply within 1 day of harvest.

POTATOES, EGGPLANT, PEPPER [illegible] Potato Beetle — 10 lbs per acre. Leafhopper, Flea Beetle, Lygus Bugs — 20 lbs per acre. Apply when insects appear. Apply dust thoroughly [illegible] and repeat application at 7 to 10 day intervals.

TOMATOES [illegible] Fruitworm, Armyworm, Flea Beetles, Tomato Hornworm, European Corn Borer, Stink Bugs — 20 lbs per acre. Apply thoroughly at first signs of insect infestation. Repeat as necessary. May be applied up to the day of harvest. Because of the [illegible] contents of this dust, it should not be used on tomatoes grown for canning or processing before clearance information has been obtained from your canner or processor.

SEVIN® SULFUR DUST 10-35

ACTIVE INGREDIENTS	[illegible]
Carbaryl (1-naphthyl N-methylcarbamate)	10.0
Sulfur	35.0
INERT INGREDIENTS	55.0
	100.0

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after using. Avoid storage near feed and food products.

Do not apply to cucurbits when [illegible] may be injurious under local conditions. Avoid [illegible] sensitive plants.

This product is highly toxic to bees exposed to direct treatment or residues on crops attractive to bees. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the area. Apply only when the hive entrance is closed. Keep bees out of treated areas for 24 hours after application. Do not apply this product or allow it to drift to blooming crops or weeds if [illegible] are present. Contact your [illegible] for protection of bees in your area.

EPA Reg. No. [illegible]-AA

[illegible]

[illegible] PUREGRO COMPANY LOS ANGELES, CALIFORNIA

NET [illegible]

LEAFY VEGETABLES AND ROOT CROPS [illegible] cabbage, Collards, Dandelion, Endive, Beet, Horseradish, Kale, Leek, Mustard Greens, Parsley, Parsnip, Radish, Rutabaga, Salsify, Spinach, Swiss chard, Turnips. Apply 10 to 20 lbs per acre to control flea beetles, cabbage [illegible] Do not apply 14 days before harvest.

ASPARAGUS: Apply 15-20 lbs per acre to control asparagus beetle. Apply during cutting season. Do not repeat application within 3 days. Do not apply 1 day before harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: Flea beetle, harlequin bug, corn earworm, armyworms. For small loopers after heads form, use SEVIN every 5 to 7 days. Apply 15-20 lbs per acre. Do not apply 3 days before harvest.

FILBERT: Filbert worm, filbert aphid, filbert leaf roller — Apply 40 lbs per acre. No time limitation.

FORAGE AND FEED CROPS

ALFALFA, CLOVERS, PEANUTS, SOYBEANS, COWPEAS: Blister beetles, Mexican bean beetle, thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar, velvetbean caterpillar and green cloverworm, armyworms, webworms, corn earworm and grasshoppers. Apply 10 lbs per acre. No time limitation.

SORGHUM: (Grain and Forage) Armyworm, webworms, corn earworm, stink bugs, sorghum midge, [illegible] direct dust into forming heads. Apply three to four days after heads have emerged from boot. Grasshoppers — Apply 20 lbs per acre. Do not apply 21 days before harvest of grain. No time limitation on sorghum forage.

RICE, FORAGE GRASSES, PASTURE: Armyworms, [illegible] per acre to control grasshoppers. Do not [illegible] to Rice within 14 days of [illegible] Pastures or [illegible]

ENVIRONMENTAL CAUTIONS

[illegible]

CONDITIONS OF SALE

[illegible]