

DIRECTIONS FOR USE

Measure the correct amount and add to the sprayer tank while it is filling with the agitator running. Stir and mix thoroughly in hand sprayers. Thorough, full coverage of spray application is important and nozzles should be adjusted properly for uniform distribution.

LETTUCE: For control of leaf-hoppers, imported cabbage worms, cabbage loopers and alfalfa loopers (California), use 2 quarts per acre in 25 to 100 gallons of water in aerial sprayers. Apply when the insects are found to be present, making thorough application. Repeat at weekly to 10 day intervals or as required for adequate insect control. Do not apply within 14 days of harvest on leaf lettuce; 7 days for head lettuce.

SPINACH: For control of leafhoppers and cabbage loopers. Use 1 quart per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial sprayers. Apply when insects are first noted, making thorough application. Repeat applications at weekly to 10 day intervals as required for adequate insect control. Do not apply within 7 days of harvest.

CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI: For control of cabbage looper and imported cabbage worm. Use 1 quart per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial sprayers. Apply when insects are first noted, making thorough applications at weekly to 10 day intervals as required for adequate insect control. Do not apply within 7 days of harvest.



MALATHION PERTHANE 4-4E

ACTIVE INGREDIENTS:

Diethyl diphenyl dichloroethane	45.0%
Related compounds	2.3%
Malathion*	41.4%
Xylene	6.3%
INERT INGREDIENTS:	5.0%
	100.0%

*O,O—DIMETHYL DITHIOPHOSPHATE OF DIETHYL MERCAPTOSUCCINATE

Contains 4 pounds Perthane per gallon.

U.S. Pat. Nos. 2,4833,428; 2,917,553

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contact with skin. In case of skin contact wash with soap and water. Avoid breathing spray mist. Avoid contamination of feed and foodstuffs. Do not take internally.

EPA REG. NO 1202-304-AA

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.




LOS ANGELES, CALIFORNIA

NET CONTENT _____ GALS.

BROCCOLI: For control of cabbage looper and imported cabbage worm. Use 1 quart of water in ground sprayers or in 5 gallons water in aerial sprayers. Apply when insects are first noted, making thorough applications at weekly to 10 day intervals as required for adequate insect control. Do not apply within 3 days of harvest.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.



This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CONDITIONS OF SALE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not separately recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.