

DIRECTIONS FOR USE

ALMONDS: For Shothole Fungus use 6 to 8 pounds per 100 gallons of water when buds begin to swell and popcorn stage.

APRICOTS: For Shothole Fungus and Brown Rot use 8 to 9 pounds per 100 gallons of water during period November 15 to December 15 or as buds begin to swell. May be used in pink bud stage for Brown Rot control.

CITRUS: For Brown Rot use 1 1/4 to 2 pounds per 100 gallons of water. Spray ground, lower branches and foliage to a height of about 3 feet from the ground. Apply 3 weeks after petal fall, also before heavy fall rains. May be combined with oil spray for scale on lemons.

CELERY: For Late and Early Blight use 6 to 9 pounds per 100 gallons of water every 10 to 14 days after celery is 6 inches high. Shorten interval between sprays during rainy and foggy weather.

PEACHES: For Leaf Curl and Shothole Fungus use 8 to 9 pounds per 100 gallons of water from November 15 to December 15 or as buds begin to swell. May be applied at pink bud stage for Brown Rot control.

PEARS AND APPLES: For Fire Blight use 1 to 1 1/2 pounds per 100 gallons of water. Do not apply more than 6 pounds per acre per application. Begin at 10 percent bloom and repeat at 5 day intervals until late bloom is over. Do not use on Comise Seckel or Anjou varieties of pears.

PLUMS: For Brown Rot use 8 to 9 pounds per 100 gallons at green bud and popcorn stage.

SOUR CHERRIES: For Brown Rot use 8 to 9 pounds per 100 gallons at pink bud, petal fall and early cover. Do not apply to sweet cherries or to English Morelo variety, as severe injury will occur.

ZN CU 19-19

When used as directed this product will aid in correcting zinc and copper deficiencies.

ACTIVE INGREDIENTS:

Zinc as metallic	19.0%
Copper as metallic	19.0%
INERT INGREDIENTS	62.0%
	100.0%

E.P.A. Reg. No. 1202-307

**CAUTION
KEEP OUT OF
REACH OF CHILDREN
MAY BE HARMFUL IF SWALLOWED**

Avoid contact with skin and eyes. Avoid breathing dust or spray mist.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

NET WEIGHT LBS

CONDITIONS OF SALE

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

ENVIRONMENTAL CAUTIONS


This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

MIXING DIRECTIONS

Place the required amount of material in the spray tank before all the water has been added. Keep agitator running constantly.

NOTE: For improved weathering ability, coverage and increased spray deposit, add 1/2 to 1 pint of PUREGRO SPREADER-DEPOSIT BUILDER. Add as the last ingredient to the spray tank.

LEAF LIFE[®]



PUREGRO
COMPANY
LOS ANGELES, CALIFORNIA