

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS
Avoid contact with skin and eyes. Avoid breathing dust or spray mist.
STATEMENT OF PRACTICAL TREATMENT
If SWALLOWED eat a physician immediately. If ON SKIN wash immediately with soap and water. If IN EYES flush eyes for at least 15 minutes. If IRRITATED remove victim to fresh air. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

10

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

"WARNING" "Area to be treated with [Product Name] on (date of application). Do not enter without appropriate protective clothing until sprays have dried. If accidentally exposed, consult the Statement of Practical Treatment portion of the pesticide label for first aid and medical attention recommendations."

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store only in original container in locked storage area. If a spill occurs, sweep up and dispose of in a permitted disposal facility. Avoid cross contamination with other pesticides and fertilizers during handling and storage. Do not store near heat or open flame. Not for Use or Storage in or Around the Home.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ZN CU 19-19

ACTIVE INGREDIENTS:

Zinc as metallic 19.0%
Copper as metallic 19.0%

INERT INGREDIENTS 62.0%

100.0%

EPA Reg No 1202-307AA
EPA Est No

KEEP OUT OF REACH OF CHILDREN

CAUTION

When used as directed this product will aid in correcting zinc and copper deficiencies.

LEAF LIFE



NET WEIGHT LBS

MIXING DIRECTIONS

1762-307
1061

Place the required amount of material in the spray tank before all the water has been added. Run agitator running constantly.

NOTE: For improved weathering ability, coverage and increased spray deposit, add 1/2 to 1 pint of PUREGRO SPREADER-DEPOSIT BUILDER. Add as the last ingredient to the spray tank.

PUREGRO CONDITIONS OF SALE AND WARRANTY

PureGro Company warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label. PureGro makes no other warranty, express or implied, of fitness or merchantability nor authorizes any agent or representative to make any variation or exception from this warranty.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of PureGro Company. Buyer assumes all risk of use, storage, or handling of this material not in strict accordance with directions given hereon.

ALMONDS: For Shothole Fungus use 6 to 8 pounds per 100 gallons of water when buds begin to swell and popcorn stage.

APRICOTS: For Shothole Fungus and Brown Rot use 8 to 9 pounds per 100 gallons of water during period November 15 to December 15 or as buds begin to swell. May be used in pink bud stage for Brown Rot control.

CITRUS: For Brown Rot use 1 1/2 to 2 pounds per 100 gallons of water. Spray ground, lower branches and foliage to a height of about 3 feet from the ground. Apply 3 weeks after petal fall, also before heavy fall rains. May be combined with oil spray for scale on lemons.

CELERY: For Late and Early Blight use 6 to 9 pounds per 100 gallons of water every 10 to 14 days after celery is 6 inches high. Shorten interval between sprays during rainy and foggy weather.

PEACHES: For Leaf Curl and Shothole Fungus use 8 to 9 pounds per 100 gallons of water from November 15 to December 15 or as buds begin to swell. May be applied at pink bud stage for Brown Rot control.

PEARS AND APPLES: For Fire Blight use 1 to 1 1/2 pounds per 100 gallons of water. Do not apply more than 6 pounds per acre per application. Begin at 10 percent bloom and repeat at 5 day intervals until late bloom is over. Do not use in Comice Seckel or Anjou varieties of pears.

PLUMS: For Brown Rot use 8 to 9 pounds per 100 gallons at green bud and popcorn stage.

SOUR CHERRIES: For Brown Rot use 8 to 9 pounds per 100 gallons at pink bud, petal fall and early cover. Do not apply to sweet cherries or to English Morelo variety as severe injury will occur.

ACCEPTED
NOV 21 1984
Under the Federal Insecticide, Fungicide, and Herbicide Act, as amended, this pesticide is registered under EPA Reg. No. 1762-307

BEST AVAILABLE COPY