

CROP	INSECT	AMOUNT/ACRE IN AT LEAST 10 GAL. WATER	DIRECTIONS
Cabbage Cauliflower Broccoli Brussels Sprouts Kale, Kohlrabi Collards	Cabbage Maggot	3 TBS/1/2 gal. water	Dip Roots: Swish roots and crowns just before setting. Water plants as soon as possible. Protection limited to newly set plants.
		2 qt./400 gal. water	Transplant drench: Apply 1/2 cup at base of each plant. Protection limited to newly set plants.
		1 qt./100 gal. water	Spray soil and newly set plants at 4, 11, and 18 days after setting. Stir or wash treatment into the soil.
	Cabbage worms Flea beetles Harlequin bug	1 qt.	Apply when insects appear. Several repeated applications may be necessary. Do not apply after edible parts begin to form.
Eggplant	Flea beetle Serpentine leaf miner	1 qt.	Make repeated applications as necessary. Do not apply after fruit begins to form.
Peppers Potatoes Sweet Potatoes	Wireworms White grubs Flea beetle larvae Cutworms	2 1/2 qt.	Work into the top few inches of soil prior to or at planting time.
Strawberry	White grubs Strawberry root weevil Strawberry rootworm	2 1/2 gal.	Work into the soil as a preplant soil treatment.
	Crown borer Spittlebugs Strawberry weevil Strawberry rootworm adult	1 1/2 qt.	Spray plants at time blossoms open or after harvest. Do not apply after fruit begins to form.
	Strawberry root weevil adult	2 qt.	Make foliage application before fruit begins to form or after harvest.
Tobacco	Wireworms	1 1/2 pt. in 300 gal. water	For transplanting water to protect transplant seedlings. Use 3 pt. for heavy infestations.
Corn (field & sweet)	Wireworms White grubs Seed corn beetles Seed corn maggots Green June bug larvae Cutworms Corn rootworms (Northern) (Southern) Corn field ants Corn root aphids	2 1/2-4 qts. broadcast	Apply as broadcast after field is plowed. Disk in promptly.
Plants African Violets	Springtails Symphilitids	5% Dust	Potting Soil—Add 1 1/2 pounds dust per gallon of soil.  Potted Plants—Sprinkle 1/4 teaspoon over soil per inch of pot. Proportionate amount for other sizes. Water from top for next several waterings. Repeat application at 4 to 6 month intervals or whenever insects appear.  Re-Potting—Mix into new soil as above. Dust old soil ball slightly.

**CAUTION:** Chlordane is toxic and can be absorbed through the skin. Do not use on pets or humans. Harmful if swallowed. Avoid inhaling spray mists. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Wash with soap and water after using. Do not use on feed or forage crops that will be fed to dairy animals or animals being fattened for slaughter. Do not apply to food crops after edible portions begin to form. Residues remaining at harvest time should be removed by washing or other effective means. Do not use on young cucurbits. Do not use on strawberries after fruit begins to form.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Apply this product only as specified on this label. Rinse equipment and containers and dispose of waste by burying in non crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with waste. Chlordane must not be used in dairy barns, milk rooms or poultry houses.

Form: ILLINUT.