

MAY 05 1989

Delta Foremost Chemical Corporation
3915 Air Park Street
Memphis, TN 38130

Attention: Haschal Golden


Gentlemen:

Subject: 1691-ES QUAD-DIS PLUS
EPA Registration Number 1203-40
Your Submission Dated December 6, 1988

The submitted final printed labels have been placed in your product file. They have not been reviewed in accordance with PF Notice 82-2.

If you have any questions concerning this letter, please contact Martha DeLaney at (703) 557-6982.

Sincerely yours,

John H. Lee 
Product Manager (31)
Antimicrobial Program Branch
Registration Division (H7505C)

54717: I: Lee: L-1: FENCO: 4/26/89: 6/7/89: KA: EK

CONCURRENCES

SYMBOL								
SURNAME								
DATE								

Delta
Foremost

1691-ES

QUAD-DIS PLUS

ACTIVE INGREDIENTS

INERT INGREDIENTS

SANITIZES DISINFECTS DEODORIZES

SOLD FOR COMMERCIAL AND INDUSTRIAL PURPOSES ONLY

DANGER: KEEP OUT OF REACH OF CHILDREN

Read carefully other precautions on right panel

Delta Foremost Chemical Corp.

DIRECTIONS
POULTRY HOUSES AND EQUIPMENT Use as directed, the product is odorless.
 (a) Remove all poultry and feed from premises.
 (b) Remove all litter and droppings from the poultry.
 (c) Empty all troughs, racks, and other feeding equipment.
 (d) Clean and disinfect thoroughly. Surfaces should be cleaned and disinfected daily.
 (e) Ventilate buildings, coops, and other equipment until treatment has been completed.
 (f) A-brested feed racks, troughs, troughs, and other equipment should be thoroughly rinsed with potable water in the hatchery to disinfect trays, incubators, and other equipment due to "exploders" and other bacteria.
FOR SANITIZATION: Clean thoroughly with a solution of concentrate and use of water with at least 10% solution.
FOR DISINFECTION: After cleaning, use a solution of concentrate and use of water with at least 10% solution.
STORAGE: Do not contaminate water, food, or feed.
PESTICIDE STORAGE AND SPILL PROCEDURE: Store in a cool, dry place. In case of exposure to extreme temperature. In case of spill, dispose of with chemical waste.
PESTICIDE DISPOSAL: Pesticide should be disposed of according to label instructions at or by an authorized agency.
CONTAINER DISPOSAL: Triple rinse for reuse, or puncture and dispose of in a sanitary manner, or by burning if burned, or by other means approved by local authorities.
PRECAUTIONS: Hazard to Humans.
KEEP OUT OF REACH OF CHILDREN. CAUTION: Irritant to eyes, on skin or on clothing. Handle with care. Wear protective gloves when handling. Avoid contamination of clothing.
STATEMENT OF WORKING CONDITIONS: In case of contact, immediately flush eyes with water. For eyes, call physician. For skin, wash with soap and water. For ingestion, drink egg whites or milk solution of it. Call a physician.
NOTE: Probable mutagenicity in experimental animals.
 Rev. 10-26-88

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

General—This product is a concentrate to be mixed with water. Do not use in conjunction with soap or other anionic detergents. It is compatible with detergents of cationic or non-ionic type.

HOSPITAL AND RESTAURANT USE

For disinfection of walls, floors, lavatory fixtures, and other hard surfaces, after cleaning, use a solution containing one ounce of concentrate per one gallon of water, and allow at least ten (10) minutes contact time.

Dishes, glassware, and other food utensils after cleaning with a solution containing one ounce of concentrate to five gallons of water at least one (1) minute contact time. Recommended by local health department recommendations.

Deodorize by stepping or spraying a solution containing one ounce of concentrate to one gallon of water.

SANITIZATION OF DAIRY EQUIPMENT

Clean and rinse thoroughly all cans, pails, buckets, and other related equipment. Then rinse with sanitizing solution containing one ounce of concentrate to five gallons of water at a least one (1) minute contact time. Drain thoroughly and air dry. The same dilution is also recommended for udder washing prior to milking. Follow recommendations of local health department.

This concentrated product fulfills the criteria of App. 10.1.1.1 of the Pasteurized Milk Ordinance 1965 Recommendations for Sanitization Service in waters up to 550 ppm (32 grains per gallon) of CaCO₃ when tested by the method outlined by the Code of Federal Regulations.

MEAT, POULTRY AND FOOD PROCESSING EQUIPMENT

Clean and rinse equipment thoroughly. Then use a solution of concentrate to five gallons of water. Allow to remain on surfaces for 10 minutes before use. On surfaces difficult to clean, use a stronger solution of concentrate to one gallon of water. Rinse thoroughly with potable water.

BEST AVAILABLE COPY