

BEST DOCUMENT AVAILABLE

ACCEPTED
MAY 16 1977
U.S. Dept. of Agriculture
1203-48

DDAP
DATE

DIRECTIONS FOR USING

Use FLY CRYSTALS where flies congregate and breed in and around such places as DAIRY BARN, MARKETS, CAFES, OPEN AIR THEATRES, POULTRY ESTABLISHMENTS, MEAT AND FOOD PROCESSING PLANTS, DRIVE-IN RESTAURANTS, ENTRANCES TO HOMES, COMMERCIAL BUILDINGS AND STORES, OUT DOOR RECREATION AREAS, such as PARKS, PICNIC GROVES, FAIRS and CAMPS, PATIOS, PORCHES, KENNELS, GARBAGE CANS, DUMPS, and INCINERATORS.

• FLY CRYSTALS GIVE YOU TWO-WAY KILL: FLY CRYSTALS are highly attractive to adult flies and kill adults that feed upon or contact the crystals if the crystals remain dry and within reach of the flies. When moisture contacts the crystals the toxicant enters the moisture and is dispersed in all directions. In this manner the poison contacts the maggots and kills them.

• FOR ADULT (Winged) FLIES: Simply scatter the crystals by hand where flies congregate. Never pile the crystals! Scatter them! For both quick and lasting reduction of flies: First, scatter the crystals over enough of the area that all the flies have an opportunity to feed upon or touch the crystals very soon; and second, also treat as many places as possible where the crystals may remain dry and uncovered. Apply 5 times weekly until the fly population is reduced. After that apply as necessary. Use 3 to 4 oz. per 100 square feet of area actually treated.

• FOR MAGGOTS: Use more care. Scatter evenly and thoroughly over the entire infested area. Use about six inches or more around all sides. Thorough, even coverage is essential.

• FOR BOTH MAGGOTS AND ADULTS: Use 1 lb. to 1 1/2 lbs. per 100 square feet of area treated. Apply to the ground floors, ledges, window sills, shelves, litter, garbage, manure and similar areas where flies congregate or breed. Apply evenly and thoroughly to breeding areas. Repeat as needed for HOUSE FLY, STABLE FLY, BLOW FLY AND MAGGOTS.

• FOR GARBAGE DUMPS, DISPOSAL AREAS: Scatter by hand or with a rotary hand duster onto the garbage or wherever the flies congregate. Use 2 to 4 oz. per 100 square feet.

• FOR USE ON THE OUTSIDE OF HOTELS, MOTELS, FOOD PROCESSING PLANTS, RESTAURANTS, SUPER-MARKETS, STOCK YARDS, POULTRY ESTABLISHMENTS: Scatter the FLY CRYSTALS thoroughly around all refuse cans, garbage and other litter where the crystals can remain dry as long as possible.

• IN LIVESTOCK BARN AND HOG PENS: Apply to dirt, manure, litter or concrete floors for an initial reduction of flies and also to as many places as available where the FLY CRYSTALS can remain dry and uncovered for a maximum length of time, such as discussed above for dairy barns.

Apply daily until fly breeding is reduced; then treatment every 3 or 4 days should be sufficient.

Delta

Foremost

4890-ES
FLY CRYSTALS

ACTIVE INGREDIENTS

1. Methionol, O, O Dimethyl phosphonate of	1.000
2. 2-methyl-4-nitrophenyl dimethyl phosphate	0.558
Other Ingredients	0.042
INERT INGREDIENTS	98.400
Total	100.000

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Read further caution statements on right panel.

NET WT. _____ LBS.

Delta Foremost Chemical Corp.

3915 AIR PARK STREET, MEMPHIS, TENNESSEE, 38118

DIRECTIONS
• USE FLY CRYSTALS on the floor of the manure area and in Entrance after floor dries. On window ledges and on Feeds. Room Flies. Leave for 24 hours after each clean up. Reapply as necessary. Sweepings to livestock. • POULTRY AND OTHER PLACES WHERE AGAINST ADULT FLIES: To one half hand in wet places heavily above for maggots. POULTRY ON GROUNDS feed upon the crystals. Maggots and adults. 12.5 lbs. per 100 sq. ft. walkways, on manure of each of the pens.

APPLICATION AREAS OF USE NOT PERMITTED

Contains DDAP which is absorbed through skin after using and may feed and foodstuffs of domestic animals. Inhibition. If per cent.

Keep tightly closed

DO NOT USE: Fly crystals should not be used on milk is processed in tanks. Finished floors under foot. To avoid rubber glove of safety. This product is not to be used on any body of water. Do not contaminate water supplies. Apply this product to waste.

The use of this product and involving electrical wiring. Expressly not for use in explicit accordance with the instructions on this label.

Do not reuse empty container by perforating or crushing.

6 27

ACCEPTED
 MAY 16 1977
 U.S. Dept. of Agriculture, Act.
 1203-48

DAIRY BARN

Delta

FOREMOST

4890-ES

FLY CRYSTALS

ACTIVE INGREDIENTS

Malathion (O, O-Dimethyl dithio-phosphate of diethyl mercaptosuccinate)	1.000
2, 2-dichlorovinyl dimethyl phosphate	0.558
Related Compounds	0.042
INERT INGREDIENTS	98.400
Total	100.000

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Read further caution statements on right panel.

NET WT. _____ LBS.

Delta Foremost Chemical Corp.

3915 AIR PARK STREET, MEMPHIS, TENNESSEE, 38118

DIRECTIONS FOR USING

USE FLY CRYSTALS IN DAIRY BARN: Scatter FLY CRYSTALS on the floor of the milking area such as Feed Chert Aisles or Stanchion area and in Entrances. For best results clean floor and apply soon after floor dries. The killing action can be prolonged by using on windows and eaves away from feed troughs, on floors not washed daily, on Feed Room floors and similar areas.

Leave for as many days as the flies continue to die or reapply after each clean-up treatment; then when fly population is reduced reapply as necessary. Do not get in feed. Do not feed the floor to livestock.

POULTRY AND RABBITS ON WIRE: Scatter on the Manure and other places where flies settle such as Feed Rooms and Egg Houses. Against adult flies, where the manure is dry, apply about one-third to one-half handful every two or three yards. Treat all damp or wet places heavily at 1 to 1 1/2 lbs. per 100 square feet as described above for maggots.

POULTRY ON GROUND OR LITTER: Apply where poultry cannot feed upon the crystals. Use in Manure Pits under Roosts against maggots and adults. Do not apply more than 1 lb. per 200 square feet (2.5 lbs. per 650 square feet). Apply in windows on ledges, in walkways on narrow shelves, and on Feed Room floors where out of reach of the poultry.

APPLICATION OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS IS NOT PERMITTED.

CAUTION

Contains DDVP and Malathion. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin; wash thoroughly after using and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep out of reach of children, poultry, and domestic animals. DDVP and Malathion may cause cholinesterase inhibition. If poisoning occurs get prompt medical aid. Atropine is antidotal.

Keep tightly closed when not in use to preserve killing power.

DO NOT USE: Fly Crystals are not to be used inside homes or where milk is processed or stored or where poultry can feed on the crystals. Finished floors may be scratched if Fly Crystals are trodden under foot. To avoid skin contact when applying Fly Crystals use a rubber glove of a type found in most drug stores.

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOTICE

The use of this product being beyond our knowledge and control and involving elements of risk if misused, we do not make any warranty expressed or implied as to the effects of any use not in explicit accordance with the claims, directions and cautions set forth on this label.

CONTAINER DISPOSAL

Do not reuse empty drums. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place from water supplies.

EPA Reg. No. 1203-48
 EPA Est. 1203-TN-1

61 277

and around
 AIR THEA-
 PROCESSING
 HOMES, COM-
 ATION AREAS,
 ODS, PATIOS,
 ICHEATORS.

CRYSTALS are
 eed upon or
 thin reach of
 xicant enters
 s manner the

stals by hand
 for them! For
 or the crystals
 opportunity to
 nd, also treat
 main dry and
 n is reduced,
 J square feet

nd thoroughly
 more around

1 1/2 lbs. per
 oors, ledges,
 similar areas
 thoroughly to
 STABLE FLY.

by hand or
 ver the flies

FOOD PRO-
 LOCK YARDS.
 S thoroughly
 the crystals

dirt, manure,
 s and also to
 s can remain
 such as dis-

ent every 3 or