

ACCEPTED
 JAN 14 1987
 1503-49

Delta

OREMOST

4891-ES
 FLY-KIL

ACTIVE INGREDIENTS:

Methomyl (S-Methyl N-((Methylcarbamoyl) Oxy) Thioacetimidate	1.127%
Z-9 Tricosene	0.026%
INERT INGREDIENTS	98.847%
TOTAL	100.000%

CAUTION

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly after using and before eating or smoking. Avoid contamination of feed and foodstuffs.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat.

IF ON SKIN: Wash thoroughly with soap and water. Call a physician if irritation persists.

IF IN EYES: Flush with plenty of water.

NOTE TO PHYSICIAN: Methomyl is a cholinesterase inhibitor. Atropine is antidotal.

Do not use this product in food areas of food handling establishments, restaurants, or other places where food is prepared or processed. Do not apply where poultry or animals, especially dogs or young calves, can pick it up or lick it.

ENVIRONMENTAL HAZARDS

Exposed surfaces treated with this product are hazardous to birds and other wildlife. Use with care when applying in areas frequented by wildlife or adjacent to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

EPA Reg. No. 1203-210

EPA Est. No. 1203-TN-1

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with labeling.

APPLICATION OF THIS PRODUCT IN THE FOOD AREA OF FOOD HANDLING ESTABLISHMENTS IS NOT PERMITTED.

DO NOT USE: Fly-Kil is not to be used inside homes or where milk is processed or stored or where poultry can feed on the crystals, inside poultry establishments, meat and food processing plants, and restaurants. Finished floors may be scratched if Fly-Kil is trodden under foot. To avoid skin contact when applying Fly-Kil, use a rubber glove of a type found in most drug stores.

Use FLY-KIL where FLIES congregate and breed in and around such places as DAIRY BARNs, MARKETs, CAFES, OPEN AIR THEATRES, ENTRANCES TO HOMES, COMMERCIAL BUILDINGS AND STORES, OUT DOOR RECREATION AREAS, such as PARKS, PICNIC GROVES, FAIRS and CAMPS, ZOOS, PATIOS, PORCHES, KENNELS, GARBAGE CANS, DUMPS, and INCINERATORS.

•**FLY-KIL GIVES YOU TWO-WAY KILL:** FLY-KIL is highly attractive to adult flies and kills adults that feed upon or contact the crystals if the crystals remain dry and within reach of the flies. When moisture contacts the crystals the toxicant enters the moisture and is dispersed in all directions. In this manner the poison contact the maggots and kills them.

•**FOR ADULT (Winged) FLIES:** Simply scatter the crystals by hand where flies congregate. Never pile the crystals! Scatter them! For both quick and lasting reduction of flies: First, scatter the crystals over enough of the area that all the flies have an opportunity to feed upon or touch the crystals very soon; and, second, also treat as many places as possible where the crystals may remain dry and uncovered. Apply 5 times weekly until the fly population is reduced. After that apply as necessary. Use 3 to 4 oz. per 100 square feet of area actually treated.

•**FOR MAGGOTS:** Use more care. Scatter evenly and thoroughly over the entire infested area. Use about six inches or more around all sides. Thorough, even coverage is essential.

•**FOR BOTH MAGGOTS AND ADULTS:** Use 1 lb. to 1½ lbs. per 100 square feet of area treated. Apply to the ground floors, ledges, window sills, shelves, litter, garbage, manure and similar areas where flies congregate or breed. Apply evenly and thoroughly, to breeding areas. Repeat as needed for HOUSE FLY, STABLE FLY, BLOW FLY AND MAGGOTS.

•**FOR GARBAGE DUMPS, DISPOSAL AREAS:** Scatter by hand or with a rotary hand duster onto the garbage or wherever the flies congregate. Use 2 to 4 oz. per 100 square feet.

•**FOR USE ON THE OUTSIDE OF HOTELS, MOTELS, FOOD PROCESSING PLANTS, RESTAURANTS, SUPER-MARKETS, STOCK YARDS, POULTRY ESTABLISHMENTS:** Scatter FLY-KIL thoroughly around all refuse cans, garbage and other litter where the crystals can remain dry as long as possible.

•**POULTRY AND RABBITS ON WIRE:** Scatter on the Manure and other places where flies settle such as Feed Rooms and Egg Houses. Against adult flies, where the manure is dry, apply about one-third to one-half handful every two or three yards. Treat all damp or wet places heavily at 1 to 1½ lbs. per 100 square feet as described above for maggots.

•**POULTRY ON GROUND OR LITTER:** Apply where poultry cannot feed upon the crystals. Use in Manure Pits under Roosts against maggots and adults. Do not apply more than 1 lb. per 200 square feet (2.5 lbs. per 650 square feet). Apply in windows on ledges, in walkways, on narrow shelves, and on Feed Room Floors where out of reach of the poultry. Keep tightly closed when not in use to preserve killing power.

STORAGE & DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place. Pesticides that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Consult federal, state or local disposal authorities for approved alternate procedures such as limited open burning.

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.