

ACCEPTED

May 24, 1967

SEARCHED INDEXED  
SERIALIZED FILED  
MAY 24 1967  
FBI - LOS ANGELES

KEEP OUT OF REACH OF CHILDREN

CAUTION

Do not breathe dust or fumes. Do not get on clothing. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contact with food and food containers. Keep out of reach of children.

Observe the following

Use Precautions

Do not use on silage forage to be fed to dairy animals or to animals intended for slaughter. Do not use on food and drug products. Do not use on feedstuffs for animals. Do not use on crops to be established. Use only at recommended rates, and do not apply closer to harvest than specified.



Agricultural Insecticide

ACTIVE INGREDIENT  
Carbaryl (1-methyl N-methylcarbamate) 100%  
INERT INGREDIENTS 10%  
100%

WARRANTY

PLANT'S WARRANTY is that this product, sold by its name, is a true and accurate representation of the quality of the product. It is not a warranty of performance or of any other quality. The user assumes all responsibility for the use of this product. The user should read the label or other instructions carefully and follow them. The user should also read the instructions on the label or other instructions. The user should also read the instructions on the label or other instructions.

Net Weight 50 lbs.

LOS ANGELES, CALIF. 90001

Apply when air is calm, or with wind equipment, at 10-15 mph. Do not apply to the ground. Do not apply to plants such as alfalfa or clover. Do not apply to the first crop of alfalfa. Do not apply to alfalfa at a rate of 10 lbs. per acre. Do not apply to alfalfa at a rate of 10 lbs. per acre.

Apply when air is calm, or with wind equipment, at 10-15 mph. Do not apply to the ground. Do not apply to plants such as alfalfa or clover. Do not apply to the first crop of alfalfa. Do not apply to alfalfa at a rate of 10 lbs. per acre. Do not apply to alfalfa at a rate of 10 lbs. per acre.

Recommendations

For control of Bollworm, Tobacco Budworm, and Cotton Leafworm: Apply 3 to 13 lbs. of Sevin 10% Dust per acre when insects first appear.  
For control of Mill and Lintworm, Bollworm, Fall Webworm, Pink Bollworm, Cotton Leafworm, Ligus Bugs, and Cotton Leaf Perforator: Apply 15 to 25 lbs. of Sevin 10% Dust per acre at 5-7 day intervals until insects first appear.

For control of Bollworm, Tobacco Budworm, and Cotton Leafworm: Apply 3 to 13 lbs. of Sevin 10% Dust per acre when insects first appear.  
For control of Mill and Lintworm, Bollworm, Fall Webworm, Pink Bollworm, Cotton Leafworm, Ligus Bugs, and Cotton Leaf Perforator: Apply 15 to 25 lbs. of Sevin 10% Dust per acre at 5-7 day intervals until insects first appear.