



**CONTROLLED IODINE
SANITIZER/CLEANER**

ECHO



Net Contents
ONE
U. S. Gallon

for **DAIRIES, MILK BARS, and all
FOOD HANDLING ESTABLISHMENTS**

DANGER
**KEEP OUT OF
REACH OF CHILDREN**

See First Aid and
Other Precautions
On Right Panel

Active Ingredients:	
Alpha-(p-Nonylphenyl)-omega-hydroxypoly (oxyethylene)-iodine complex*	16.75%
Phosphoric Acid	10.00%
Inert Ingredients	73.25%
*Providing 1.75% Titratable Iodine	

EPA Registration No. 1214-31
EPA Est. No. 1214-GA-1

ECHO PRODUCTS
MANUFACTURER • ATLANTA, GA. 30301

E-05.3-P

• Rinse empty container thoroughly with water and discard it. •

WHAT IODOX IS... AND DOES

Iodox is a fast acting sanitizer which kills most pathogenic and common food spoiling micro-organisms. It is considered non-selective in its action and is not adversely affected by cold water or water hardness. Iodox cleans, sanitizes and prevents the formation of milkstone. The rich amber color of Iodox in use dilutions is its own indicator of germicidal activity - as long as the color remains, Iodox is germicidally effective.

DIRECTIONS

For cleaning and sanitizing areas where food is prepared:

For non-porous surfaces: Scrub with a solution of 3 oz. Iodox in 5 gallons of water.
• For porous surfaces or those difficult to clean: Scrub with a solution of 6 oz. Iodox in 5 gallons of water. • A terminal rinse with potable water is required on all food contact areas.

For removing milkstone:

Use 1 1/2 oz. Iodox in 3 gallons of water. • Rinse surfaces with potable water before use.

For sanitizing pre-cleaned utensils, glassware, dishes, etc.:

Use a solution of 1 oz. Iodox in 5 gallons of water (25 PPM) and immerse pre-cleaned utensils, glassware, dishes into the solution for at least two minutes. Drain and allow to dry.

DANGER:

Causes eye damage and skin irritation. • Do not get in eyes, on skin, or on clothing. • Protect eyes and skin when handling. • Harmful if swallowed.
• Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. • For eyes, call a physician. • Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. • Call a physician immediately.

This product fulfills the criteria of Appendix F of the Grade 'A' Pasteurized Milk Ordinance 1965 Recommendations of the U. S. Public Health Service