


INSECT	AMOUNT TO USE	RECOMMENDATION
TERMITE CONTROL	In 4 gal. water 1 gal.	Remove all wood from contact with soil. Treat along foundation wall of buildings with deep foundations by digging trenches 30 inches deep and 8 inches wide. Do not dig below the footing. Use 4 gal. to every 5 linear feet of trench. Apply half in trench, balance to soil as it is replaced. For treating infestation under concrete porches and other inaccessible areas, drill holes near house foundation at 2 feet intervals and pour in 1 gal. Inspect annually for reinfestation.
LAWN INSECTS Chiggers Lawn Moths (Sod Webworms) Ants Digger Wasps White Grubs Earth Worms Wireworms Cutworms	In 5 gal. water to 1000 sq. ft. 1/2 pt.	Apply to entire area at start of season as an overall treatment. Wash in with water after 48 hours. Repeat if needed for chiggers and lawn moths, using one-fourth pint.
HOUSEHOLD INSECTS	In 1 gal. water	Apply thoroughly with coarse spray or brush to cracks, surfaces and other areas where insects are seen resting, running or hiding. Repeat as necessary. Do not use in bedrooms.
Roaches, Waterbugs	1/3 pt.	Treat areas where roaches hide.
Ants	1/3 pt.	Treat ant trails and areas around foundation, door sills, window frames where ants crawl.
Silverfish, Firebrats	1/3 pt.	Treat baseboards, areas around shelving, bookcases and storage areas.
Carpet Beetles	1/3 pt.	Apply to infested area of floor, baseboard and underside of carpet.
Brown Dog Tick In premises	1/3 pt.	Treat infested cracks, hiding places, also sleeping quarters of pets. Do not treat animals. Provide fresh bedding after treatment.
Spiders, Centipedes In dwelling	1/3 pt.	Apply only where spiders or webs or centipedes are found.
Aid in control of Flies and Mosquitoes	1/3 pt.	For outside use only. Apply to resting places such as porch walls, ceilings, around windows, doors and screens.
Wasps	1/3 pt.	Spray nests in evenings. Remove and destroy nests afterward.
Crickets	1/3 pt.	Treat closet floors, other hiding places.
SHADE TREES, ORNAMENTALS	In 100 gal. water	
Bagworms	1 qt.	Apply in late spring when insects are first observed.
Leaf Beetles, Thrips Elm Leaf Beetles Willow Leaf Beetles	1 qt.	Apply to entire plant when insects are first seen.
Black Vine Weevil	1 qt.	Treat entire plant. Drench ground under the plant and branches closest to ground. Apply in late June.
Mosquitoes	4 1/2 tsp. per gal. to 1000 sq. ft.	Apply to shrubs, grass, and mosquito breeding areas. Will kill fish.



Net Contents 1 Gal.

CHLORDANE

Emulsion Concentrate

For Control of: Termites
and Certain other insects

CAUTION

KEEP OUT OF REACH OF CHILDREN

See remainder of CAUTION statement on side of panel.

Contains: 4 lbs. of Technical Chlordane per gallon. Emulsifies readily with water.

Active Ingredients: 90%

Technical Chlordane* 45%

Petroleum Hydrocarbons 45%

Inactive Ingredients: 10%

*Equivalent to 27% Octachloro-4,7-Methano-1,2,3,4,5,6-hexahydroindane and 18% Related Compounds.

USDA Reg. 1222-51

Distributed by

INDIANA FARM BUREAU COOPERATIVE ASSN., INC.

INDIANAPOLIS, INDIANA

ACCEPTED

1/14/51

1222-51

CROP	INSECT	AMOUNT IN AT 10 GAL
Cabbage Cauliflower Broccoli Brussels Sprouts Kale, Kohlrabi Collards	Cabbage Maggot	3 TBS/1/2
	Cabbage worms Flea beetles Harlequin bug	2 qt./400
Eggplant	Flea beetle Serpentine leaf miner	1 qt./100
Peppers Potatoes Sweet Potatoes	Wireworms White grubs Flea beetle larvae Cutworms	1 qt.
Strawberry	White grubs Strawberry root weevil Strawberry rootworm	2 1/2 qt.
	Corn borer Spittlebugs Strawberry weevil Strawberry rootworm adult	1 1/2 qt.
	Strawberry root weevil adult	2 qt.
Tobacco	Wireworms	1 1/2 pt. in water
Corn (field & sweet)	Wireworms White grubs Seed corn beetles Seed corn maggots Green Jung bug larvae Cutworms Corn rootworms (Northern) (Southern) Corn field ants Corn root aphids	2 1/2-4 qts.
Plants African Violets	Springtails Symphylids	5% Dust

CAUTION: Chlordane is toxic and can be absorbed through the skin. Avoid inhaling spray mists. Avoid contact with food. Wash with soap and water after using. Do not use on animals being fattened for slaughter. Do not apply to remaining at harvest time should be removed by washing. Do not use on strawberries after fruit begins to form.

This product is toxic to fish and wildlife. Keep out of water. Apply this product only as specified on this label. Do not bury in non crop lands away from water supplies. Do not bury with waste. Chlordane must not be used in

INSECT	AMOUNT TO USE	RECOMMENDATION
TERMITE CONTROL	In 44 gal. water 1 gal.	Remove all wood from contact with soil. Treat along foundation wall of buildings with deep foundations by digging trenches 30 inches deep and 8 inches wide. Do not dig below the footing. Use 4 gal. to every 5 linear feet of trench. Apply half in trench, balance to soil as it is replaced. For treating infestation under concrete porches and other inaccessible areas, drill holes near house foundation at 2 feet intervals and pour in 1 gal. Inspect annually for reinfestation.
LAWN INSECTS Chiggers Lawn Moths (Sod Webworms) Ants Digger Wasps White Grubs Earth Worms Wireworms Cutworms	In 5 gal. water to 1000 sq. ft. 1/2 pt.	Apply to entire area at start of season as an overall treatment. Wash in with water after 48 hours. Repeat if needed for chiggers and lawn moths, using one-fourth pint. Keep children and pets off treated area until grass is washed off and dried.
HOUSEHOLD INSECTS	In 1 gal. water	Apply thoroughly with coarse spray or brush to cracks, surfaces and other areas where insects are seen resting, running or hiding. Repeat as necessary. Do not use in bedrooms.
Roaches, Waterbugs	1/3 pt.	Treat areas where roaches hide.
Ants	1/3 pt.	Treat ant trails and areas around foundation, door sills, window frames where ants crawl.
Silverfish, Firebrats	1/3 pt.	Treat baseboards, areas around shelving, bookcases and storage areas.
Carpet Beetles	1/3 pt.	Apply to infested area of floor, baseboard and underside of carpet.
Brown Dog Tick in premises	1/3 pt.	Treat infested cracks, hiding places, also sleeping quarters of pets. Do not treat animals. Provide fresh bedding after treatment.
Spiders, Centipedes in dwelling	1/3 pt.	Apply only where spiders or webs or centipedes are found.
Aid in control of Flies and Mosquitoes	1/3 pt.	For outside use only. Apply to resting places such as porch walls, ceilings, around windows, doors and screens.
Wasps	1/3 pt.	Spray nests in evenings. Remove and destroy nests afterward.
Crickets	1/3 pt.	Treat closet floors, other hiding places.
SHADE TREES, ORNAMENTALS	In 100 gal. water	
Bagworms	1 qt.	Apply in late spring when insects are first observed.
Leaf Beetles, Thrips Elm Leaf Beetles Willow Leaf Beetles	1 qt.	Apply to entire plant when insects are first seen.
Black Vine Weevil	1 qt.	Treat entire plant. Drench ground under the plant and branches closest to ground. Apply in late June.
Mosquitoes	4 1/2 tsp. per gal. to 1000 sq. ft.	Apply to shrubs, grass, and mosquito breeding areas. Will kill fish.



Net Contents 1 Gal.

CHLORDANE

Emulsion Concentrate

For Control of: Termites
and Certain other insects

CAUTION

KEEP OUT OF REACH OF CHILDREN

See remainder of CAUTION statement on side of panel.

Contains: 4 lbs. of Technical Chlordane per gallon. Emulsifies readily with water.

Active Ingredients: 90%

Technical Chlordane* 45%

Petroleum Hydrocarbons 45%

Inactive Ingredients: 10%

*Equivalent to 27% Octachloro-4,7-Methano-Tetrahydroindane and 18% Related Compounds.

USDA Reg. 1222-51

Distributed by

INDIANA FARM BUREAU COOPERATIVE ASSN., INC.

INDIANAPOLIS, INDIANA

CROP	INSECT	AMOUNT IN 10 GAL.
Cabbage Cauliflower Broccoli Brussels Sprouts Kale Kohlrab Collards	Cabbage Maggot	1 qt.
	Cabbage worms Flea beetles Harlequin bug Leaf miner	1 qt.
Peppers Potatoes Sweet Potatoes	Wireworms White grubs Flea beetle larvae Cutworms	1 qt.
Strawberry	White grubs Strawberry root weevil Strawberry rootworm	1 qt.
	Crown borer Spittlebugs Strawberry weevil Strawberry rootworm adult	2 qt.
	Strawberry root weevil adult	2 qt.
Tobacco	Wireworms	1 1/2 pt. in 3 water
Corn (field & sweet)	Wireworms White grubs Seed corn beetles Seed corn maggots Green June bug larvae Cutworms Corn rootworms (Northern) (Southern) Corn field ants Corn root aphids	2 1/2-4 qts. b
Plants African Violets	Springtails Symphylids	5% Dust

CAUTION: Chlordane is toxic and can be absorbed thru swallowed. Avoid inhaling spray mists. Avoid contamination. Wash with soap and water after using. Do not use on animals being fattened for slaughter. Do not apply to remaining at harvest time should be removed by washing. Do not use on strawberries after fruit begins to form.

This product is toxic to fish and wildlife. Keep out of a killed. Apply this product only as specified on this label, burying in non crop lands away from water supplies. Ca and burying with waste. Chlordane must not be used in d

DATION

ect with soil. Treat along
with deep foundations by
deep and 8 inches wide.
g. Use 4 gal. to every 5
half in trench, balance to
ing infestation under con-
cessible areas, drill holes
et intervals and pour in
infestation.

of season as an overall
or after 48 hours. Repeat
lawn moths, using one-

reated area until grass is

spray or brush to cracks,
ere insects are seen rest-
at as necessary. Do not

ide.

around foundation, door
nts crawl.

ound shelving, bookcases

or, baseboard and under-

g places, also sleeping
at animals. Provide fresh

webs or centipedes are

to resting places such as
indows, doors and screens.

move and destroy nests

g places.

sects are first observed.

n insects are first seen.

ground under the plant
nd. Apply in late June.

and mosquito breeding



Net Contents 1 Gal.

CHLORDANE Emulsion Concentrate

For Control of: Termites
and Certain other insects

CAUTION

KEEP OUT OF REACH OF CHILDREN

See remainder of CAUTION statement on side of panel.

Contains: 4 lbs. of Technical Chlordane per
gallon. Emulsifies readily with water.

Active Ingredients: 90%

Technical Chlordane* 45%

Petroleum Hydrocarbons 45%

Inactive Ingredients: 10%

*Equivalent to 27% Octachloro-4,7-Methano-
Tetrahydroindane and 18% Related Compounds.

USDA Reg. 1222-51

Distributed by

INDIANA FARM BUREAU COOPERATIVE ASSN., INC.

INDIANAPOLIS, INDIANA

CROP	INSECT	AMOUNT/ACRE IN AT LEAST 10 GAL. WATER	DIRECTIONS
Cabbage Cauliflower Broccoli Brussels Sprouts Kale, Kohlrabi Collards	Cabbage Maggot	3 TBS 1/2 gal. water	Dip Roots: Swish roots and crowns just before setting. Water plants as soon as possible. Protection limited to newly set plants.
		2 qt. 400 gal. water	Transplant drench. Apply 1/2 cup at base of each plant. Protection limited to newly set plants.
		1 qt. 100 gal. water	Spray soil and newly set plants at 4, 11, and 18 days after setting. Stir or wash treatment into the soil.
	Cabbage worms Flea beetles Harlequin bug	1 qt.	Apply when insects appear. Several repeated applications may be necessary. Do not apply after edible parts begin to form.
Eggplant	Flea beetle Serpentine leaf miner	1 qt.	Make repeated applications as necessary. Do not apply after fruit begins to form.
Peppers Potatoes Sweet Potatoes	Wireworms White grubs Flea beetle larvae Cutworms	2 1/2 qt.	Work into the top few inches of soil prior to or at planting time.
Strawberry	White grubs Strawberry root weevil Strawberry rootworm	2 1/2 gal.	Work into the soil as a preplant soil treat- ment.
	Crown borer Spittlebugs Strawberry weevil Strawberry rootworm adult	1 1/2 qt.	Spray plants at time blossoms open or after harvest. Do not apply after fruit begins to form.
	Strawberry root weevil adult	2 qt.	Make foliage application before fruit begins to form or after harvest.
Tobacco	Wireworms	1 1/2 pt. in 300 gal. water	For transplanting water to protect transplant seedlings. Use 3 pt. for heavy infestations.
Corn (field & sweet)	Wireworms White grubs Seed corn beetles Seed corn maggots Green June bug larvae Cutworms Corn rootworms (Northern) (Southern) Corn field ants Corn root aphids	2 1/2 4 qts. broadcast	Apply as broadcast after field is plowed. Disk in promptly.
Plants African Violets	Springtails Symphilitids	5% Dust	Potting Soil—Add 1 1/2 pounds dust per gallon of soil. Potted Plants—Sprinkle 1/4 teaspoon over soil per inch of pot. Proportionate amount for other sizes. Water from top for next several waterings. Repeat application at 4 to 6 month intervals or whenever insects appear. Re-Potting—Mix into new soil as above. Dust old soil ball slightly.

CAUTION: Chlordane is toxic and can be absorbed through the skin. Do not use on pets or humans. Harmful if
swallowed. Avoid inhaling spray mists. Avoid contamination of feed and foodstuffs. Keep out of reach of chil-
dren. Wash with soap and water after using. Do not use on feed or forage crops that will be fed to dairy animals
or animals being fattened for slaughter. Do not apply to food crops after edible portions begin to form. Residues
remaining at harvest time should be removed by washing or other effective means. Do not use on young cucurbits.
Do not use on strawberries after fruit begins to form.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be
killed. Apply this product only as specified on this label. Rinse equipment and containers and dispose of waste by
burying in non crop lands away from water supplies. Containers should be disposed of by punching holes in them
and burying with waste. Chlordane must not be used in dairy barns, milk rooms or poultry houses.