



For Use on	To Control	In 100 Gal.	In 1 Gal.	Last Application (days) before harvest.	Comments
Flowers Ornamentals	Oyster Shell Scale	1 pt.	1 tsp.		Apply in summer when infestation occurs. Repeat in 10 to 12 days. For scales, apply when crawlers have settled on foliage. Injury may occur on certain ferns and on some species of Crassula.
	Lacebugs				
	Aphids, Mites, White Fly, Thrips, Mealy Bug, Japanese Beetle Adult, Four Lined Plant Bug, Rose Leafhopper, Potato Leafhopper, European Pine Shoot Moth, Scurfy Scale, Eucymus Scale, Tarnished Plant Bug	1 1/2 pt.	1 1/2 tsp.		
	Begonia, Juniper Scale, Tent Caterpillar, Magnolia Scale	2 pt.	2 tsp.		
	Soft Brown Scale	2 1/2 pt.	2 1/2 tsp.		
	Pine Needle Scale	4 pt.	4 tsp.		
Tomatoes	Russet Mites	1/2 pt.	1 1/2 tsp.	3	Repeat as needed.
	Aphids	1 pt.	1 tsp.		
Beans	Mexican Bean Beetle	1 1/2 pt.	1 1/2 tsp.	3	
	Mites, Leafhopper				
Peas	Aphids	1 1/2 pt.	1 1/2 tsp.	3	
Potatoes	Aphids	1 pt.	1 tsp.	3	
Cabbage, Kohlrabi, Broccoli*, Kale, Mustard, Turnips	Imported Cabbage Worm, Cabbage Looper	1-2 pt.	2 tsp.	7	*Do not apply to broccoli within 3 days of harvest.
Eggplants	Aphids, Mites	1 pt.	1 tsp.	3	Repeat as needed.
	Lacebug	3 pt.	3 tsp.		
Peppers	Aphids	1 pt.	1 tsp.	3	
Melons, Cucumbers, Pumpkins, Squash	Aphids, Mites	1 1/2 pt.	1 1/2 tsp.	3	Make full coverage spray. Do not use when plants are wet.
Spinach, Endive, Carrot, Radish, Parsnip, Fennel, Horse-Radish, Cellar, Swiss Chard, Parsley	Aphids	2 pt.	2 tsp.	7	Repeat as needed.
Lettuce	Aphids, Mites, Leafhoppers	2 pt.	2 tsp.	7	Repeat as needed.
Onions	Thrips	1 1/2 pt.	1 1/2 tsp.	3	Repeat as needed.
Asparagus	Asparagus Beetle	*2 pt. per acre		3	*In sufficient water for good coverage. Repeat as needed.
Corn	Sap Beetle, Corn Earworm, Thrips	1 1/2 pt. per acre		5	Repeat as needed. Apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals as infestation warrants. Re-apply immediately after rain.
Barley, Oats, Rye, Wheat	Greenbugs, English Grain Aphids, Army Worms	1 1/2 pt. per acre, 2 pt.		7	Repeat as needed.
Tobacco	Mites, Aphids	1 1/2 pt.	1 1/2 tsp.	7	Repeat as needed.
Alfalfa	Aphids, Leafhoppers	1 1/2 pt. per acre in at least 10 gal. water		7	Repeat as needed.
Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries	Mites, Thrips, Leafhoppers, Japanese Beetles	1 1/2 pt.	2 tsp.	7	Make thorough coverage. Repeat as needed.
Peanut, Soybean	Aphids, Spider Mites, Leafhoppers	1 1/2 pt. per acre	1 1/2 t.	7	Repeat as needed.



Net Content: 1 Gal.

Malathion

EMULSIFIABLE CONCENTRATE

For Control of:

FLIES AND CERTAIN PLANT PESTS

CAUTION

KEEP OUT OF REACH OF CHILDREN
See remainder of caution statement on side of panel.

ACTIVE INGREDIENTS:

Malathion*	By Weight	57%
Xylene		35%

INERT INGREDIENTS: 8%

*0,0-dimethyl dithiophosphate of diethyl mercapto succinate

Contains 5 lbs. Malathion per gallon


Distributed by

INDIANA FARM BUREAU CO-OP
INDIANAPOLIS, IND.

FARM BUREAU

EPA Reg. 1222-52

For Use on Homes
 To Control: Roaches, Spiders, Ants, Crickets, Clover Mites, Silverfish, Carpet Beetles, Clothes Moths Larvae
 In 100 Gal.: 4 gal.
 In 1 Gal.: 1/3 pt.
 Comments: Apply where insects are seen. For Carpet Beetles and Clothes Moths, apply to baseboards, floors, (including areas under and along margins of carpets and in closets) infested surfaces of carpeting, other lint accumulation areas.
 *As a household in premise treatment.



Net Content: 1 Gal.

Orthion

ORTHION
 For Control of:

CERTAIN PLANT PESTS

OF CHILDREN

on statement on side of panel.

By Weight
 57%
 35%
 8%

of diethyl mercapto succinate
 Malathion per gallon
 tributed by

M BUREAU CO-OP
APOLIS, IND.

Reg. 1222-52

For Use on	To Control	In 100 Gal.	In 1 Gal.	Comments
<u>Homes</u>	Roaches, Spiders Ants, Crickets Clover Mites Silverfish Carpet Beetles Clothes Moths Larvae	4 gal.	1/3 pt.	Apply where insects are seen. For Carpet Beetles and Clothes Moths, apply to baseboards, floors, (including areas under and along margins of carpets and in closets) infested surfaces of carpeting, other lint accumulation areas.
<u>Chickens</u> <u>Turkeys</u> <u>Ducks, Geese</u>	Poultry Lice Northern Fowl Mite Chicken Red Mite*	7 pt.	2 tbs.	*As a supplement to premise treatment, apply 1 gal. to 100-150 birds.
<u>Poultry Houses</u>	Poultry Lice Northern Fowl Mite Chicken Red Mite Flies	1 1/4 gal.	4 tbs.	Apply liberally to walls, ceilings, roosts, nests and adjacent areas. Force spray into cracks and crevices. Repeat as needed. Use 1 pt. per 150 ft. of roost.
<u>Swine</u>	Lice	1 gal.	2 1/2 tbs.	Apply a complete coverage. Repeat if needed.
<u>Beef Cattle</u>	Ticks Lice	1-2 gal. 1 gal.	3-5 tbs. 3 tbs.	Apply a complete coverage spray. Repeat if needed. Do not make accessible to lactating dairy cows or calves under 1 mo. of age.
<u>Back Rubber</u>	Lice Reduction Horn Fly Reduction	9 tbs. in 1 gal. fuel oil		
<u>Barns, Hoghouses</u> <u>Poultry Houses,</u> <u>Manure Piles,</u> <u>Garbage Cans</u>	Aid in controlling House Flies Stable Flies	2 gal.	1/6 pt.	Add 1/2 cup sugar per gal. Apply where flies rest. Combine with a good sanitation program.

CAUTION: Do not contaminate feed, food, water or containers. Keep out of reach of children. Do not use in milk processing rooms. Repeat applications as needed.

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before reuse. Destroy containers completely when empty, do not reuse. Malathion may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal.

This product is toxic to fish. Keep out of lakes, streams and ponds. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by breaking and burying with wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CAUTION: DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.