

## DIRECTIONS FOR USE

CROP	INSECTS	RECOMMENDATIONS
Lawns	Ants, Chiggers Digger Wasps Sod Webworms (Lawn Moths) White Grubs, Earthworms Wireworms	Treat entire area at start of season, applying 5 lbs. per 1000 sq. ft. In case of mound building ants, work into top few inches of loose soil. Wash in with water for soil insects after 48 hours. Repeat if needed for chiggers and lawn moths. <del>Keep children and pets off treated areas.</del> Birds feeding on treated areas may be killed.
Cucumbers Melons Squash Pumpkins	Squash Bugs Striped Cucumber Beetle Spotted Cucumber Beetle	Apply 20 to 40 lbs. per acre, making first application in seedling stage and subsequent applications as insects appear. Do not apply after edible portions begin to form. Observe plants after first application as some varieties may be injured by repeated applications.
<del>Alfalfa</del> Soybeans <del>Clovers</del>	Grasshoppers, Crickets Tarnished Plant Bug Stink Bug	Apply at the rate of 30 lbs. to 40 lbs. per acre when insects first appear and repeat when necessary. Do not apply to forage crops to be fed to dairy animals or to animals being finished for slaughter, or to be sold commercially or shipped interstate.
Soil Insects under shade trees, ornamentals, perennial and annual flowers	Wireworms, White Grubs Green June Beetle Japanese Beetle, Sowbugs Millepedes	Apply 50 to 100 lbs. per acre to surface of soil, and disk in prior to seeding or transplanting. User is cautioned that heavy applications may be detrimental to plant growth. <u>For small areas apply 1 lb to 435 sq. ft.</u>
Onions, Glads Cabbage Cauliflower Brussels Sprouts Broccoli	Thrips, Cabbage Looper Imported Cabbage Worm Diamond Back Moth  Cabbage Maggots	Apply 20 to 40 lbs. per acre, making application as soon as insects appear, making 1 to 4 applications at 10 day intervals. Do not use after heads or other edible parts of plants begin to form. Do not apply to green or spring onions.  Dust plants and soil around plants at <del>4, 11 and 18 days after setting.</del> Apply at rate of 40 lbs. per acre.
	Ants, House Spiders Roaches (Waterbugs) Silverfish, Crickets Centipedes	Make thorough treatments where insects are found, dusting lightly in cracks, crevices and other surfaces frequented by insects. Do not use in bedrooms. Repeat as necessary.

### CAUTION:

<p>Avoid inhalation of dust. Keep out of reach of children.</p> <p>Avoid contamination of feed and foodstuffs.</p> <p>Chlordane is toxic to pollenizing insects, so do not apply while plants are blooming.</p> <p>This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label.</p> <p>Empty containers should be disposed by burying with waste or by burning (keep out of smoke). Avoid spillage.</p>	<p>Do not use on leafy vegetables as spinach, lettuce and celery.</p> <p>May injure certain varieties of cucurbits.</p> <p><del>Do not use on crops within 30 days of harvest.</del> <del>Remove any residues at harvest by washing or brushing.</del></p>
---	--

Form LNLHDI

Net Weight 1 Lb.

USDA-REG. NO. 1222-66

preplanting, planting time or transplanting time.

**DUST NO. 14**

**5% CHLORDANE DUST**

For the Control of

**Ants, Chiggers, Roaches  
Silverfish**

and certain other insects  
Containing 5% by weight of  
technical chlordane

FARM  
BUREAU



INGREDIENTS:

ACTIVE INGREDIENTS—  
Technical Chlordane\* . . . . . 5%

INERT INGREDIENTS . . . . . 95%

\*Equivalent to 3% Octachloro-4,7-methanotetrahydroindane and 2% of related compounds.

Note: 20 lbs. of Co-op Dust No. 14 contains an equivalent of 1 lb. of technical chlordane.

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**  
See remainder of CAUTION statement on side of panel.

Manufactured for

**INDIANA FARM BUREAU  
COOPERATIVE ASSN., INC.**

Indianapolis, Indiana