

Net Weight - 50 lbs.

ROL 7.76 PAK

(Contains Rabon Oral Larvicide)

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle and House Flies in the manure of treated swine

ACTIVE INGREDIENT	BY WEIGHT
2-Chloro-1,4,4,5-tetrachlorophenyl vinyl dimethyl phosphate	7.76%
INERT INGREDIENTS	92.24%
TOTAL	100.00%

Refers only to ingredients which are not toxic.
 RABON insecticide contains 35 grams of RABON per pound.
 RABON is the registered trademark of Shell Oil Company.
 U.S. Patent No. 3,102,842

KEEP OUT OF REACH OF CHILDREN CAUTION

See below for additional precautionary statements

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

1222-114

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 - Cattle containing this product may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.
 - Start feeding RABON larvicide feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.
 - When fed, this product passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch.
 - It prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle and house flies in treated swine, but is not effective against existing adult flies.
 - In some cases, supplementary control measures may be needed in and around cattle pens and barns to control adult house flies and stable flies which can breed not only in manure but in other decaying vegetable matter or sewage on the premises.
 - In order to achieve optimum fly control, this product should be used in conjunction with effective good management and sanitation practices.
 - This product will mix uniformly in feeds when standard mixing procedures are followed. Thus usual precautions that are common to a feed preparation and which cause stratification such as excessive free-fall or excessive handling are to be avoided. It is recommended that appropriate pre-feeding techniques be employed to assure adequate distribution throughout the feed mix. The premix should be pre-mixed with other feed ingredients in a separate container and then mixed with the feed. Mixing should be adequate to assure uniform dispersion. Optimum performance can be assured only if this product is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixers, horizontal blenders, mixer-feeders, trucks) may be used to prepare formulated feeds.
 - Feeds prepared with this product should not be heated, nor be mixed with feeds containing predominantly pellets. Further, this product should not be mixed in liquid feed supplements.
- CATTLE - In a Concentrate Feed-Roughage Fed Separately**
- This product can be used to prepare concentrate feeds that will provide 792 mg of RABON per animal per day.
 - To prepare a larvicide concentrate feed, mix this product according to the amount of concentrate fed per animal per day. Use the following table as a guide for determining the proper mix rate.

Pounds of Concentrate Consumed per animal per day	MIXING DIRECTIONS FOR CATTLE RABON in the Concentrate		Pounds of this product per ton of Concentrate
	mg/lb	%	
.5	15.4	0.35	900
1.0	79.2	0.18	450
1.5	528	0.12	300
2.0	396	0.07	225
5.0	159	0.03	90
10.0	79	0.018	45
15.0	53	0.012	30
20.0	39	0.009	23
25.0	32	0.007	18
30.0	26	0.006	15

* Feed the appropriate larvicide concentrate to cattle weighing between 400 and 1200 pounds. For larger cattle weighing between 1200-1700 pounds, increase the amount of premix per ton of concentrate to 1.5 times that recommended.

CATTLE - In a Complete Ration-No Other Roughage Fed

- This product can be used to prepare rations that contain 26.4 mg of RABON per pound of complete ration.
- To prepare a larvicide ration, mix 1.5 pounds of this product per ton of complete ration containing both grain and roughage.
- For feeds this larvicide complete ration to feeder cattle weighing from 400 to 1400 pounds or to dairy cattle at a rate to sustain milk production, but not less than 2.6 pounds of the ration per 100 pounds of body weight daily.

SWINE - All swine should be treated

- Pigs (weaners to market weight)
- Mix 1.5 lbs. of this product per ton of meal type feed and offer free choice. This is equivalent to 22.7 mg of RABON per lb. of feed.
- Sows (lactating and breeding gilts)
- Mix 2.6 lbs. of this product per ton of meal type feed and offer 4 lbs. of feed per animal per day. This is equivalent to 45.4 mg of RABON per lb. of feed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Pesticide that cannot be reused should be disposed of according to applicable Federal, State or Local procedures.
 Container Disposal: Containers must be disposed of properly. Do not reuse or rent containers. Empty containers should be disposed of in a sanitary manner. Do not reuse or rent containers.

Manufactured By

INDIANA FARM BUREAU
COOPERATIVE ASS'N., INC.

Main Office: Indianapolis, IN 46204

EPA Reg. No.
EPA Est. No.

<p align="center">U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDE PROGRAMS REGISTRATION DIVISION, WASHINGTON, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input checked="" type="checkbox"/> REREGISTRATION <i>Under the Federal Insecticide, Fungicide, and Rodenticide Act.</i></p>	EPA REGISTRATION NO.	DATE OF ISSUANCE
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT	
<p>NAME AND ADDRESS OF REGISTRANT (include ZIP code)</p> <div style="text-align: right; padding-right: 50px;"> <p>DRP 1/4/53</p> <p>160 83294 15 1</p> </div>		
<p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>		
<p><input type="checkbox"/> ATTACHMENT IS APPLICABLE</p>		
SIGNATURE OF APPROVING OFFICIAL		DATE

f. The producing EPA Establishment No. should appear on the label in the appropriate location.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George F. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (FS-767)

Enclosure