

MANUFACTURED BY

D/S/D-NP

FOR INDUSTRIAL USE ONLY

DISINFECTANT/SANITIZER/DEODORIZER STAPHYLOCIDAL/SALMONELLICIDAL/PSEUDOMONICIDAL

nomes.	
USE DILUTION TEST AOAC:	
Staphylococcus aureus ATCC #6538	1.85
Salmonella cholerasuis ATCC #10708	
Pseudomonas aeruginosa ATCC #15442	
	1.00
Fungicidal against pathogenic fungi Trichophyton interdigitale	4.05
Strain #640	1:85
ACTIVE INGREDIENTS:	
n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18)	
dimethyl benzyl ammonium chlorides	3.00%
n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl	
ammonium chlorides	3.00%
Sodium Carbonate	
Tetrasodium ethylene diamine tetraacetate	
	11.00%
INERT INGREDIENTS	89.00%
D/S/D-NP does not contain phosphorous compounds	

DANGER: KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR FIRST AID AND OTHER PRECAUTIONS

EPA REG. NO. 1239-36

Rinse empty container thoroughly with water and discard

DIRECTIONS:

SANITIZING: In non-hospital applications, use $\frac{1}{2}$ oz. of D/S/D-NP to each gallon of water for cleaning and sanitizing of inanimate surface areas. Laboratory tests have shown D/S/D-NP will reduce the bacteria level by 99.9%.

DISINFECTING: Use $1\frac{1}{2}$ oz. of D/S/D-NP to each gallon of water for cleaning and disinfecting of inanimate surface areas. It has been proven that D/S/D-NP is Staphylocidal, Salmonellicidal, and Pseudomonicidal. It is fungicidal against Trichophyton interdigitale, the causative organism of Athlete's Foot.

When using D/S/D-NP on food contact areas, surfaces must be rinsed with potable water prior to reuse. D/S/D-NP may be applied to floors with the use of an autoscrubber or mop. A vacuum pickup may be used, but do not rinse with fresh water. Apply D/S/D-NP at 1½ oz. per gallon of water to clean, deodorize and disinfect walls, furniture bedframes and bedpans. Apply the D/S/D-NP solution with a cloth, mop or sponge.

DANGER: KEEP OUT OF REACH OF CHILDREN.

CORROSIVE. Causes eye damage and severe skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food. Not for use or storage in or around the home.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.