

PARATHION 2% DUST

ACTIVE INGREDIENTS:

Parathion*	2%
INERT INGREDIENTS:	98%
Total	100%

*O, O-diethyl O-p-nitrophenyl phosphorothioate

DANGER

KEEP OUT OF REACH OF CHILDREN

May be Fatal If Swallowed, Inhaled, or Absorbed Through Skin! Rapidly Absorbed Through Skin. Do not get in eyes or on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.



ANTIDOTE AND FIRST AID TREATMENT:

INTERNAL: If swallowed give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY.** (Note to Physician— Therapeutic doses of atropine appear effective as an antidote.)

EXTERNAL: If skin is contacted, remove contaminated clothing and immediately wash skin with soap and water.

USE PRECAUTIONS

Toxic to fish and wildlife. Do not apply in any manner not specified on the label. Do not contaminate any body of water or apply when conditions favor drift or runoff from the crop area treated.

Rinse equipment and dispose of wastes by burying in non-crop land away from water supplies. Empty bags should be disposed of by burying with wastes or burning. (KEEP OUT OF SMOKE).

DIRECTIONS FOR USE

LETTUCE, CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, COLLARDS, KALE, SPINACH, MUSTARD GREENS, TURNIPS and ONIONS: For control of aphids, thrips, cabbage looper and cabbage worms apply 20 to 25 lbs. per acre when insects appear in damaging numbers. Do not apply within 21 days of harvest of lettuce, or within 15 days of harvest of spinach or onions, or within 10 days of harvest of other crops.

CANTALOUPE and WATERMELONS: For control of spider mites, leafhoppers, thrips, aphids and leafminer apply 20 to 25 lbs. per acre when insects appear in damaging numbers. Do not apply within 7 days of harvest.

LEGUMES, such as ALFALFA, CLOVER, VETCH and on small grains such as BARLEY, OATS, and WHEAT: For control of aphids, leafhoppers, armyworms, blister beetles and grasshoppers apply 12 to 20 lbs. per acre depending on infestation. Do not use within 15 days of harvest.

PEAS and TOMATOES: For control of aphids apply 15 to 20 lbs. per acre. Repeat application 7-10 days later if needed. Do not apply within 10 days of harvest.

BEANS and POTATOES: For control of aphids, red spider, mexican bean beetle, Colorado potato beetle and leafminer apply 15 to 20 lbs. per acre, repeating applications at 7-10 day intervals if insects persist. Do not apply within 15 days of harvest of beans or within 5 days of harvest of potatoes.

CORN: For control of European corn borer, grasshopper and armyworm apply 20 to 25 lbs. per acre. On corn borer consult your State Agricultural Experiment station as to proper timing of applications. Four applications at 5-day intervals beginning with the egg hatch is often advised, especially for sweet corn. Do not make application within 12 days of picking, cutting or forage use.

GRAPES: For control of mealybug, leaf folder, and red spider apply 20 to 25 lbs. per acre, repeating at 5-7 day intervals for red spider. Do not apply within 14 days of harvest.

STRAWBERRIES: For control of aphids and red spider apply 15 to 20 lbs. per acre, repeating against red spider 7-10 days later. Do not apply within 14 days of harvest.

COTTON: For control of aphids, red spider mites, fleahopper and garden webworms apply 15 to 20 lbs. per acre. Do not apply within 5 days of hand picking or within 15 days if cotton trash is to be fed to dairy animals.

TOBACCO: See label — For control of aphids and midge (suck fly), green June beetle grubs: dust bed thoroughly as needed. Field — For control of aphids and grasshoppers apply 12 to 15 lbs. per acre. Do not apply within 7 days of priming or 15 days of cutting. Handlers should wear gloves. Avoid getting wet with plant juices when cutting.

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damage arising from a breach of this warranty.

LR198E-10R 6023841
USDA Reg. No. 1258-106

SAVAGE, MINN
LEBANON, PA
HOUSTON, TEX
NO. LITTLE ROCK, ARK
OMAHA, NEB

AGRICULTURAL DIVISION



LITTLE ROCK, ARKANSAS

O L I N CHEMICAL CORPORATION

PHOENIX, ARIZ
RICHMOND, IND
ST. LOUIS, MO
AUGUSTA, GA
WICHITA, KAN

Manufactured by
**ZEP MANUFACTURING
COMPANY**
ATLANTA, GEORGIA 30301

Division of
National Service Industries, Inc.

U.S. Pat. 3,111,539; 3,111,539

ZEP TOXII

PRESSURIZED SPRAY

KILLS WASPS,
SPIDERS, HORNETS,
BEES AND
YELLOW JACKETS

CAUTION

May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing of spray mist and provide adequate ventilation of area being treated. Do not smoke while using. Contact with skin, eyes, or clothing should also be avoided. Wash thoroughly with soap and warm water after handling. If poisoning occurs, get prompt medical aid.

To Physician—Atropine sulfate is antidotal.

Avoid contamination of food, utensils, food preparation areas. Avoid contamination of fish ponds. Remove pets from area before spraying.

Avoid excessive wetting of plastic, rubber and asphalt surfaces.

Do not use in commercial food preparation areas or in edible products areas of food processing plants.

WARNING!

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 120° F. may cause bursting. Never throw container into fire or incinerator.

EPA REG NO 1270-19/AA
EPA EST NO 1270 GA 1



Effective at a distance of 12 feet



**DOUBLE ACTION —
KILLS WASPS AND OTHER
INSECT PESTS BY FREEZING
AND INSECTICIDAL ACTION
FOR OUTDOOR USE ONLY.**

ACTIVE INGREDIENTS:

Pyrethrins	.10%
*Piperonyl butoxide, technical	.20%
†N-octyl bicycloheptene dicarboximide	.33%
**2-(1-Methylethoxy)phenol methylcarbamate	.50%
Petroleum distillate	.84%
INERT INGREDIENTS	98.03%
Total 100.00%	

*Consists of .16% (butylcarbityl) (6-propylpiperonyl) ether and .04% other related compounds.
†Baygon® U. S. Pat. No. 3,111,539.
‡MGK 264, Insecticide Synergist.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Contents under pressure. See side panel for additional precautionary statements

NET WEIGHT 1 lb. 2 oz. AV.

READ DIRECTIONS CAREFULLY

Point spray nozzle toward insect's nest, hole or crevice, where infestation is known to exist. Spray area until saturated with material. Also spray around in the general area until it is completely wet. This treated area will have a residual killing action and prevent reinfestation for 2 to 4 weeks. If the residual film becomes ineffective on account of rain or other causes, repeat treatment.

Always spray with the wind at the user's back
DIELECTRIC

Spray Test: Electrodes 11" apart; no arcing at 35,000 volts.

Cup Test: Electrodes 1/10" apart; no breakdown up to 25,000 volts.

WASPS ARE KNOCKED DOWN BY:

1. Lowering of body temperature
2. Suffocation
3. Paralysis

Cotton States 5% Sevin[®] Garden Dust

U. S. D. Reg. No. 1339-159

DIRECTIONS

APPLICATION: Begin applications when insects or damage is first observed and repeat on a regular 7 to 10-day schedule or as necessary for good insect control.

VEGETABLES

BEANS: To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, and Corn Earworm—apply at the rate of 20 pounds per acre or 1 ounce per 50 feet of row. Where Mexican Bean Beetle is the only problem, a rate of 10 pounds per acre is sufficient. To control Lygus Bugs, Stink Bug and Lima Bean Pod Borer—apply 40 pounds per acre. This product can be applied up to the day of harvest.

CUCUMBER, SUMMER SQUASH: To control Pickleworm and Melonworm—apply at the rate of 10 pounds per acre. To control Cucumber Beetles, Squash Bugs, and Flea Beetles—apply at the rate of 20 pounds per acre or 1 ounce per 50 feet of row. This product can be applied up to the day of harvest.

TOMATO, EGGPLANT, PEPPER: To control Tomato Fruitworm, Fall Armyworm, Flea Beetles and Leafhoppers—apply at the rate of 20 pounds per acre or 1 ounce to 50 feet of row. For control of Tomato Hornworm, European Corn Borer and Stink Bugs—apply at the rate of 40 to 80 pounds per acre. This product can be applied up to the day of harvest.

POTATO: To control Colorado Potato Beetle—apply at the rate of 10 pounds per acre. For control of Leafhoppers and Flea Beetles apply at 20 pounds and for Lygus Bugs apply at 40 pounds per acre.

ACTIVE INGREDIENTS
Carbaryl—1-Naphthyl N-Methylcarbamate 5.0%
INERT INGREDIENTS 95.0%
TOTAL 100.0%
[®] Trademark of Union Carbide Corp. for Carbaryl—1-Naphthyl N-Methylcarbamate.

MANUFACTURED BY



MONROE, LOUISIANA

ORNAMENTALS

To control Apple Aphid, Bagworm, Birch Leaf Miner, Blister Beetles, Boxelder Bug, Boxwood Leaf Miner, Elm Leaf Aphids, Elm Leaf Beetle, Flea Beetles, Japanese Beetle, Lace Bugs, Leafhoppers, Leaf Rollers, Oak Leaf Miners, Orange Tortrix, Periodical Cicada, Plant Bugs, Psyllids, Rose-slug, Rose Aphids, Scale insects in the crawler stage, Tent Caterpillars, Willow Leaf Beetle and exposed Thrips—apply thoroughly to the infested plants at first sign of damage and repeat at 7-day intervals or as necessary for good insect control. To obtain scale control apply in Spring and early Summer when scale crawlers are present.

LAWNS

To control Leafhoppers, Fall Armyworm, and Sod Webworm—apply at the rate of 4 pounds per 1,000 square feet. For best results, apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 to 3 weeks if necessary.

CAUTION


Keep out of Reach of Children

Harmful if swallowed or inhaled. Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing of dust. SKIN CONTACT MAY BE HARMFUL. Avoid contact. Wash hands before eating. Take shower or bath after work. WEAR REGULAR LONG SLEEVED WORK CLOTHING. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is Antidotal.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.



1339-159