

NET
CONTENTS



GALLONS

HERBICIDE

BUTYL ESTER D6 WEED KILLER

ACTIVE INGREDIENT:

Butyl Ester of 2,4-Dichlorophenoxyacetic Acid
(Equivalent to 2,4-Dichlorophenoxyacetic Acid 63.0%)

79.1%

INERT INGREDIENTS:

TOTAL

20.9%

100.0%

Contains the equivalent of 6 lbs. of 2,4-Dichlorophenoxyacetic Acid per gallon at 68° F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION. Avoid contact with eyes, skin and clothing. Do not take internally. Do not contaminate feed or foodstuffs.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Course sprays are less likely to drift.) Do not contaminate domestic or irrigation water supplies. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not mix with liquid fertilizers.

Because of the difficulties of removing 2,4-D and 2,4,5-T from sprayers do not use the same spray equipment for applying other materials to plants as injury will result. Use a separate sprayer for this purpose.

DO NOT STORE NEAR FERTILIZERS, SEEDS, INSECTICIDES, FUNGICIDES OR PLANTS

GENERAL INFORMATION

If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using. When filling spray tank first fill tank 1/2 full of water; add the calculated amount of 2,4-D emulsifiable; then finish filling tank with water with by-pass open and pump running; if oil is used as carrier, follow same procedure and stir thoroughly. Material may tend to settle when not agitated. When mixture is left in sprayer allow by-pass pressure to re-agitate before resuming spraying.

PRECAUTION

Do not reuse empty container. Destroy by burying in a safe place. Do not burn. This material is toxic to fish. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label.

TIME OF APPLICATION

Annual weeds are most susceptible in the seedling stage and early growth stages of development and when conditions are such as to promote a vigorous growth. WHEN USED IN CROP SPRAYING AS A SELECTIVE SPRAY, THE STAGE OF GROWTH OF THE CROP MUST ALSO BE CONSIDERED, OTHERWISE SEVERE DAMAGE MAY RESULT.

SOME SUSCEPTIBLE WEEDS

Arrowweed	Henbit	Puncture Vine
Antelope	Horseshoe	Redtop
Beggar Ticks	Jimsonweed	Redtop
Bull Thistle	Kochia	Shepherd's Purse
Buttercup	Lamb Quarters	Spartan Weed
Catnip	Loco Weed	St. John's Wort (annual)
Chickory	Mallow	Stinkweed
Cocklebur	Mastard	Succas
Dandelion	Mexican Weed	Sunflower
Dock	Mustard Glory	Wild Mustard
Fameweed	Mustard	Wild Onion
Figwort	Nettle	Wild Parsnip
Four O'Clock	Pepper Grass	Wild Radish
Ground Ivy	Pig Weed	Wild Sweet Potato
Wild Rape	Prickly Lettuce	

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damage arising from a breach of this warranty.

LR147G-03U 6224371
EPA Reg. No. 125B-921

Use drip nozzles when crop is 10 inches tall or higher. Keep spray off leaves. Do not spray after rainfall appears. Some injury to corn may occur. Do not apply during hot or dry stages. Do not spray or graze treated grain fields within two weeks after treatment. Do not feed treated straw to livestock.

When crop is small and more resistant to 2,4-D but when weeds are in sensitive growth stage. In spring when weeds are small and grain is in full tiller stage (4 to 8 inches high) and always before boot stage. After plants are fully tillered but before stems begin to joint. When plants have formed 4 or 5 leaves but before plants have come into bud or bloom stage.

1/4 lb.	1/3 pt.	1/5 pt.
FALL	1/3 - 2/3 pt.	1/4 - 1/3 pt.
SPRING	1/3 - 1/2 pt.	1/4 - 1/3 pt.
1/2 lb.	1/2 pt.	1/5 pt.

CORN, MILO, WHEAT, OATS, BARLEY, RYE, FLAX

Some varieties of corn should not be sprayed (Solex, Serrano, Royal, Kona, De la Cruz, etc.). Do not overdo. Consult your State agricultural experiment station for recommendations under local conditions or more specific uses.

Manufactured For Olin

125 F-221