

Crop	*Amount D-6 Per Acre	Equivalent 2,4-D Per Acre	APPLICATION	CAUTION
CORN, MILO	1/3 pt.	1/4 lb.	When crop is small and more resistant to 2,4-D but when weeds are in sensitive growth stage.	Use drop nozzles when crop is 10 inches tall or higher. Keep spray off leaves. Do not spray after tassels appear. Some injury to corn may occur.
WHEAT OATS BARLEY RYE	FALL 1/3 - 2/3 pt. SPRING 1/3 - 1/2 pt.	FALL 1/4 - 1/2 lb. SPRING 1/4 - 1/3 lb.	In spring when weeds are small and grain is in full tiller stage (4 to 8 inches high) and always before boot stage. After plants are fully tillered but before stems begin to joint.	Do not apply during boot or dough stage. Do not forage or graze treated grain fields within two weeks after treatment. Do not feed treated straw to livestock.
FLAX	1/5 pt.	1/7 lb.	When plants have formed 4 or 5 leaves but before plants have come into bud or bloom stage.	Some varieties should not be sprayed (tolerant varieties: Redwing, Sheyenne, Royal, Kota, Dakota.) Do not overdose.

*Application rates may vary slightly from State to State depending upon soil types and weather factors. Consult your State agricultural experiment station for recommendations under local conditions or more specific uses.

NET
CONTENTS



GALLONS

BUTYL ESTER D6 WEED KILLER

ACTIVE INGREDIENT:

Butyl Ester of 2,4-Dichlorophenoxyacetic Acid
(Equivalent to 2,4-Dichlorophenoxyacetic Acid 63.0%)

79.1%

INERT INGREDIENTS:

TOTAL

20.9%

100.0%

Contains the equivalent of 6 lbs. of 2,4-Dichlorophenoxyacetic Acid
per gallon at 68° F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION. Avoid contact with eyes, skin and clothing. Do not take internally. Do not contaminate feed or foodstuffs.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Course sprays are less likely to drift.)

Do not contaminate domestic or irrigating water supplies.

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not mix with liquid fertilizers.

Because of the difficulties of removing 2,4-D and 2,4,5-T from sprayers do not use the same spray equipment for applying other materials to plants as injury will result. Use a separate sprayer for this purpose.

DO NOT STORE NEAR FERTILIZERS, SEEDS, INSECTICIDES, FUNGICIDES
OR PLANTS

GENERAL INFORMATION

If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using. When filling spray tank first fill tank 1/4 full of water; add the calculated amount of 2,4-D emulsifiable, then finish filling tank with water with by-pass open and pump running. If oil is used as carrier, follow same procedure and stir thoroughly. Material may tend to settle when not agitated. When mixture is left in sprayer allow by pass pressure to re-agitate before resuming spraying.

PRECAUTION

Do not reuse empty container. Destroy by burying in a safe place. Do not burn. This material is toxic to fish. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label.

TIME OF APPLICATION

Annual weeds are most susceptible in the seedling stage and early growth stages of development and when conditions are such as to promote a vigorous growth. WHEN USED IN CROP SPRAYING AS A SELECTIVE SPRAY, THE STAGE OF GROWTH OF THE CROP MUST ALSO BE CONSIDERED, OTHERWISE SEVERE DAMAGE MAY RESULT.

SOME SUSCEPTIBLE WEEDS

Arrowhead	Henbit	Puncture Vine
Artichoke	Horsetail	Purslane
Beggar Ticks	Jimsonweed	Ragweed
Bull Thistle	Kochia	Shepherd's Purse
Butterprint	Lambs Quarters	Smart Weed
Catnip	Loco Weed	Sow Thistle, (annual)
Chickory	Mallow	Stinkweed
Cocklebur	Marestail	Sumacs
Dandelion	Mexican Weed	Sunflower
Docks	Morning Glory	Wild Mustard
Fanweed	Mustards	Wild Onion
Figwort	Nettles	Wild Parsnip
Four O'Clock	Pepper Grass	Wild Radish
Ground Ivy	Pig Weed	Wild Sweet Potato
Wild Rape	Prickly Lettuce	

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damage arising from a breach of this warranty.

LR147G-03U 6224371
USDA Reg. No. 1258-221

Manufactured For
AGRICULTURAL DIVISION **Olin**
LITTLE ROCK, ARKANSAS
OLIN CORPORATION

CROP	*Amount D4 Per Acre	Equivalent 2,4-D Per Acre	APPLICATION	CAUTION
CORN, MILO	1/2 pt.	1/4 lb.	When crop is small and more resistant to 2,4-D but when weeds are in sensitive growth stage.	Use drop nozzles when crop is 10 inches tall or higher. Keep spray off leaves. Do not spray after tassels appear. Some injury to corn may occur.
WHEAT OATS BARLEY RYE	FALL 1/2 - 1 pt. SPRING 1/2 - 2 1/2 pt.	1/4 - 1/2 lb. 1/4 - 1/2 lb.	In spring when weeds are small and grain is in full tiller stage (4 to 8 inches high) and always before boot stage. After plants are fully tillered but before stems begin to joint.	Do not apply during boot or dough stage. Do not spray or graze treated grain fields within two weeks after treatment. Do not feed treated straw to livestock.
FLAX	1/4 pt.	1/8 lb.	When plants have formed 4 or 5 leaves but before plants have come into bud or bloom stage.	Some varieties should not be sprayed (tolerant varieties: Redwing, Sheyenne, Royal, Kata, Dakota.) Do not overdose.

*Application rates may vary slightly from State to State depending upon soil types and weather factors. Consult your State agricultural experiment station for recommendations under local conditions for more specific uses.



HERBICIDE

BUTYL ESTER ¹²⁵⁸

WEED **D4** KILLER ^{6-14-68 1258-722}

ACTIVE INGREDIENT:

Butyl Ester of 2,4-Dichlorophenoxyacetic Acid 58.8%

INERT INGREDIENTS:

TOTAL 41.2%

2,4-Dichlorophenoxyacetic Acid Equivalent 46.9%

Contains the equivalent of 4 lbs. of 2,4-Dichlorophenoxyacetic Acid per gallon at 68° F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION. Avoid contact with eyes, skin and clothing. Do not take internally. Do not contaminate feed or foodstuffs.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Course sprays are less likely to drift.)

Do not contaminate domestic or irrigating water supplies.

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not seed lawns, gardens or fields until at least four weeks after application of spray to area. Do not mix with liquid fertilizers.

BECAUSE OF THE DIFFICULTIES OF REMOVING 2,4-D and 2,4,5-T FROM SPRAYERS DO NOT USE THE SAME SPRAY EQUIPMENT FOR APPLYING OTHER MATERIALS TO PLANTS AS INJURY WILL RESULT. USE A SEPARATE SPRAYER FOR THIS PURPOSE.

DO NOT STORE NEAR FERTILIZERS, SEEDS, INSECTICIDES, FUNGICIDES OR PLANTS

AVAGLE, MINN.
CHAS. J. PA
MICHIGAN, ILL.
E. L. BROWN, ILL.
GAYLOR, N.E.H.

AGRICULTURAL DIVISION **Olin**
E. L. BROWN, ILL.



HERBICIDE

BUTYL ESTER 1258

WEED D4 KILLER 6-14-68 1258-722

ACTIVE INGREDIENT:

Butyl Ester of 2,4-Dichlorophenoxyacetic Acid 58.8%

INERT INGREDIENTS:

TOTAL 41.2%

100.0%

2,4-Dichlorophenoxyacetic Acid Equivalent 46.9%

Contains the equivalent of 4 lbs. of 2,4-Dichlorophenoxyacetic Acid per gallon at 68° F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION. Avoid contact with eyes, skin and clothing. Do not take internally. Do not contaminate feed or foodstuffs.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Course sprays are less likely to drift.)

Do not contaminate domestic or irrigating water supplies.

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not seed lawns, gardens or fields until at least four weeks after application of spray to area. Do not mix with liquid fertilizers.

BECAUSE OF THE DIFFICULTIES OF REMOVING 2,4-D and 2,4,5-T FROM SPRAYERS DO NOT USE THE SAME SPRAY EQUIPMENT FOR APPLYING OTHER MATERIALS TO PLANTS AS INJURY WILL RESULT. USE A SEPARATE SPRAYER FOR THIS PURPOSE.

DO NOT STORE NEAR FERTILIZERS, SEEDS, INSECTICIDES, FUNGICIDES OR PLANTS

GENERAL INFORMATION

If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using. When filling spray tank first fill tank 1/4 full of water; add the calculated amount of 2,4-D emulsifiable, then finish filling tank with water with by-pass open and pump running. If oil is used as carrier, follow same procedure and stir thoroughly. Material may tend to settle when not agitated. When mixture is left in sprayer allow by-pass pressure to re-agitate before resuming spraying.

PRECAUTION

Do not reuse empty container. Destroy by burying in a safe place. Do not burn. This material is toxic to fish. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label.

TIME OF APPLICATION

Annual weeds are most susceptible in the seedling stage, and early growth stages of development and when conditions are such as to promote a vigorous growth. WHEN USED IN CROP SPRAYING AS A SELECTIVE SPRAY, THE STAGE OF GROWTH OF THE CROP MUST ALSO BE CONSIDERED, OTHERWISE SEVERE DAMAGE MAY RESULT.

SOME SUSCEPTIBLE WEEDS

Arrowhead	Henbit	Puncture Vine
Artichoke	Horseail	Pursh
Beggar Ticks	Jimsonweed	Ragweed
Bull Thistle	Kochia	Shepherd's Purse
Butterprint	Lamb's Quarters	Smart Weed
Catnip	Loco Weed	Sow Thistle, (annual)
Chickory	Mallow	Stinkweed
Cocklebur	Marestail	Sumac
Dandelion	Mexican Weed	Sunflower
Docks	Morning Glory	Wild Mustard
Famweed	Mustards	Wild Onion
Figwort	Nuttall	Wild Parsnip
Four O'Clock	Pepper Grass	Wild Radish
Ground Ivy	Pig Weed	Wild Sweet Potato
Wild Rape	Prickly Lettuce	

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damage arising from a breach of this warranty.

LR267AG1-04R 6224337
USDA Reg. No. 1258-222

AGRICULTURAL DIVISION **Olin**

CHEMICAL CORPORATION

PHOENIX, ARIZ.
CHICAGO, ILL.
ST. LOUIS, MO.
ATLANTA, GA.
WICHITA, KAN.