

1258-279

03/21/1956

1/2

With letter FEB 23 1956

CONTENTS

GALLONS

CONTROL NO.

MATHIESON BRAND TRADE MARK

TERRACLO[®] 2 lb. EMULSIFIABLE CONCENTRATE

A fungicide for the control of certain soil-borne diseases.

COMPOSITION

ACTIVE INGREDIENT:

Pentachloronitrobenzene ----- 24%

INERT INGREDIENTS: ----- 76%

TOTAL ----- 100%

Contains 2 lbs. of Pentachloronitrobenzene per U.S. gallon at 70°F.

CAUTION

Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face thoroughly after using. Wear gloves and respirator when loading the spreader or sprayer.

Do not use or store near heat or open flame. Use adequate ventilation. Keep container closed.

DIRECTIONS FOR USE

Root and Stem Rot of Green Beans caused by *Rhizoctonia solani*:

Apply 2 to 3 3/4 gal. of this product per acre in 10 gals. of water. Spray at 30 to 40 P.S.I. pressure in a band six to twelve inches wide over the seed and surrounding soil at planting time. TERRACLO[®] was developed for this use in Florida.

Sclerotinia Leaf Drop of Lettuce:

Apply 7 1/2 to 11 1/4 gal. of this product per acre in 150 to 200 gals. of water. The spray should be applied to wet the lettuce and the soil surface in a band six to twelve inches wide, centered on the row. Application is made when plants are two to three inches high and have been thinned.

Bottom Rot of Lettuce:

Make two applications each consisting of 2 gal. of this product per acre in 150 to 200 gals. of water. The spray should be applied so as to wet the lettuce and the soil surface in a band six to twelve inches wide, centered on the row. First treatment should be applied when plants are two to three inches high, second treatment ten days thereafter.

Stem Rot of Carnations, African Violets, Poinsettias, Chrysanthemums, Snapdragons, Begonias and similar plants attacked by *Rhizoctonia solani*:

As a soil drench apply this product at the rate of 7 1/2 pints in 300 gallons of water per 1000 square feet of bench or bed area. For best results, apply to a well prepared seed bed.

ACCEPTED

MAR 21 1956

1258-279

With letter **MAR 19 1956**

With letter **FEB 23 1956**

2/2

TERRACLOR* 2 lb. EMULSIFIABLE CONCENTRATE

Crown Rot (*Sclerotinia trifoliorum*) of Clovers and Alfalfa:

Use 4 1/2 to 6 gals. of this product in 100 gallons of water per acre, apply spray on the soil surface within one week after cutting. For best control, use in the late fall or when the first signs of disease development are apparent.

Damping-off (*Rhizoctonia solani*) of Cotton:

Use 1 1/2 gallons per acre of ~~TERRACLOR 2 lb. A.C.~~ ^{this product} mixed with at least 15 gallons of water. Apply as a spray upon the seed and surrounding soil at planting time.

IMPORTANT NOTICE: Since the use of this product is beyond the control of the seller, seller makes no warranty of any kind, express or implied, concerning the use of this product. User assumes all risk of use or handling whether in accordance with directions or not.

*Terraclor is a trade mark of Olin Mathieson Chemical Corporation

OLIN MATHIESON CHEMICAL CORPORATION

Baltimore, Maryland

Little Rock, Arkansas

47107

1258-279

ACCEPTED

