

1258-279

01/31/1958
with letter JAN 21 1958

1/1

ADDITIONAL DIRECTIONS FOR USE FOR Terraclor 2 lb. E. C.

BEANS (Root and stem rot - Rhizoctonia solani):

Band: Apply 2 to 3 3/4 gallons of this product per acre in 10 gallons of water. Spray at 30 to 40 P.S.I. pressure in a band 6 to 12 inches wide over the seed and surrounding soil at planting time.

Furrow: Apply 3/4 to 1 gallon of this product per acre in 15 to 20 gallons of water. Spray the planting furrow and covering soil at planting time.

NOTE: Do not feed treated vines to livestock.

BEANS - SNAP AND DRY (White mold - Sclerotinia sclerotiorum):

Apply 1 gallon of this product in 15 to 20 gallons of water per acre in an 8 inch band, centered on the row, immediately after or at the time of seeding. (Where dinitro type pre-emergence herbicides are used, add 1 gallon to the final herbicide spray mix and apply simultaneously.) If disease problem is severe, application may be repeated at two to three week intervals, using one or two nozzles per row directing spray at base of plant. Do not apply after first bloom. Soil should remain undisturbed after treatment. Terraclor was developed for this use in New York.

NOTE: Do not feed treated vines to livestock.

WHEAT (Common Smut or Bunt):

As a slurry treatment, apply in conventional slurry treatment equipment at the following rates:

For 23 cc cup, use a mixture of 1/2 gallon per gallon of water.

For 46 cc cup, use a mixture of 1 quart per gallon of water.

This will deposit approximately 1/2 ounce active per bushel of seed. Treated seed should be labeled:

"Treated seed - do not use for feed or food purposes."

ACCEPTED
1-31-58
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTRATION
ED UNDER NO. 1258-279