

Olin

Terraclor®

2 LB. EMULSIFIABLE CONCENTRATE
Soil Fungicide

Controls certain soil-borne diseases of
Cotton • Beans • Garlic • Potatoes • Lawns • Ornamentals

ACTIVE INGREDIENT:
Pentachloronitrobenzene 24.0%
INERT INGREDIENTS: 76.0%
Total 100.0%

Contains 2 lbs. of Pentachloronitrobenzene per U. S. gallon at 68° F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face thoroughly after using.

Do not store near heat or open flame. Use with adequate ventilation. Keep container closed. Avoid contamination of feed and foodstuffs.

CAUTION COMBUSTIBLE LIQUID

DO NOT USE, POUR, SPILL OR STORE
NEAR HEAT OR OPEN FLAME.

Terraclor and Olin

are OLIN Trademarks

AGRICULTURAL DIVISION
LITTLE ROCK, ARKANSAS

OLIN CORPORATION

Olin

USE CAUTIONS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not plant any root crop not listed on this label in rotation with PCNB treated soils.

DESTROY THIS CONTAINER WHEN EMPTY - NEVER RE-USE

DIRECTIONS FOR USE

BEANS (Root and Stem Rot - *Rhizoctonia solani*): FURROW - Application rates are to 14,500 linear feet of row for bush type beans and to 8,430 linear feet of row for pole beans. Apply 2-3 qts. in 8 to 10 gallons of water. Spray planting furrow and covering soil at planting time. Avoid applying directly to seed; otherwise delayed emergence may occur. In California and Arizona apply 3-4 pints in 8 to 10 gals. of water. Apply in-furrow and incorporate with covering soil at planting. Use lower rate for lighter soils. Avoid application to bare seed.

SNAP and DRY BEANS (White Mold-Sclerotinia sclerotiorum): Use 1 gal. mixed in 8 to 10 gals. of water and apply to 14,500 linear feet of row for bush type beans or to 8,430 linear feet of row for pole beans. Spray in an 8" wide band, centered on the row, immediately after or at time of seeding. Avoid applying directly to seed; otherwise delayed emergence may occur. (Where dinitro type pre-emergence herbicides are used, add 1 gallon to the final herbicide spray mix and apply simultaneously). If disease is severe, application may be repeated at 2 to 3 week intervals, using 1 or 2 nozzles per row and directing spray at base of plants. Do not apply after first bloom. Soil should remain undisturbed after treatment. Do not feed treated vines to livestock.

COTTON (Damping Off - *Rhizoctonia solani*): Use 1/2 to 1 gallon per acre in at least 15 gallons of water (per 13,000 linear feet of row based on 40" spacings). Spray on seed and surrounding soil at planting time. Do not allow the feeding or grazing of cotton foliage by livestock.

GARLIC (White Rot - *Sclerotium cepivorum*): Use 10 gals. mixed in 100 gals. of water and apply to 21,800 linear feet of row at planting time. Attach a suitable rig to the machine planter, using 2 nozzles per planting furrow. Direct the front nozzle to spray the bottom of the open furrow and the "seed" as it is dropped. Direct the rear nozzle to spray the covering soil.

POTATOES (*Rhizoctonia solani*) - BROADCAST APPLICATION: - Apply 9 to 12 1/2 gallons mixed with 100 to 150 gallons of water to one acre. Apply to the soil surface, prior to planting and disc and cross disc to a depth of 4 to 6 inches. A drag harrow is usually attached behind the discs on the last discing. It is very important to thoroughly mix the Terraclor into the tuberforming zone of the soil. It is desirable to make treatment when the soil is slightly moist.

DIRECTIONS FOR USE (cont.)

POTATOES (*Rhizoctonia solani*) - ROW (in furrow) APPLICATION: Use 5 gals. per 12,400 linear feet of row mixed in sufficient water to total approximately 20 gallons of finished spray. Spray several points in the planting row to treat all the soil that surrounds and covers the seed piece and potato forming zone. Three nozzles attached to the planter work best; one fan nozzle ahead of opening disc (approximately 5 gals. of spray per acre); one cone nozzle placed above and slightly ahead of closing discs (approximately 10 gals. of spray per acre); one fan nozzle to rear of closing discs (approximately 5 gals. of spray per acre). Spray pattern should treat 12 to 14 inch band centered on the row.

ORNAMENTALS

AZALEA AND CAMELLIA (Azalea Petal blight and Camellia flower blight): Apply as a drench under plants at a rate of 1 1/2 qts. in 30 gallons of water per 150 sq. ft. Begin application prior to opening buds and repeat every 3 to 4 weeks through the bloom period.

LAWNS (Brown Patch - *Rhizoctonia solani*): For Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses. Make one application in the fall or spring at first indication of infection. Mix 1 qt. in sufficient water (10-15 gals.) to cover 1,000 sq. ft. Apply by means of a sprinkler can, spray or any other convenient method of obtaining even coverage. Following application, the treated area should be lightly watered to insure movement through the grass to the soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3-4 weeks later. Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

For Dichondra: Treat as above, using 1 1/2 qts. mixed in 40 gals. of water per 1,000 sq. ft.

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

LR370G-04U 6324963 USDA Reg. No. 1258-279-AA

1258-279

06/14/1971

