

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions, not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

WARNING

Do not use or store near heat or open flame. During Commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator approved by the U.S. Department of Agriculture for Endrin protection. Keep unprotected humans and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until the drifting insecticides and volatile residues have dissipated.

PRECAUTIONS IN USING

Highly toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift of spray from the crop area being treated.

DESTROY THIS CONTAINER WHEN EMPTY — Puncture and crush beyond any possible re-use. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies.

USDA Reg. No. 1258-443

LRL94F-06R

SAVAGE, MINN.
LEBANON, PA.
HOUSTON, TEX.
NO. LITTLE ROCK, ARK.
OMAHA, NEB.

ENDRIN

1.6 L.B.

EMULSIFIABLE

ACTIVE INGREDIENTS:

Endrin* 19.7%
Methylated aromatic petroleum derivatives 75.4%

INERT INGREDIENTS:

Total 4.9%

100.0%

*Hexachlorocyclopentadiene, octahydro-endro, endo-dimethanonaphthalene
Contains 1.6 lbs. Endrin per U. S. gallon at 68° F.

RED

DANGER

KEEP OUT OF REACH OF CHILDREN

POISONOUS BY SKIN CONTACT, INHALATION OR SWALLOWING, RAPIDLY ABSORBED THROUGH SKIN. SPRAY MIST EXTREMELY HAZARDOUS. Do not get in eyes, on skin or on clothing. Do not breathe spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Clean up spillage. Do not use indoors without adequate exhaust ventilation. Use clean clothing daily and wear clean synthetic rubber gloves. In case of contact immediately remove all contaminated clothing and wash skin or eyes with plenty of water for at least 15 minutes; for eyes, get medical attention. Wash clothing before re-use. Avoid contamination of feed or foodstuffs.



POISON

ANTIDOTE: INTERNAL: If any of the insecticide is swallowed; give a tablespoon of salt in a glass of warm water. Repeat until vomit fluid is clear. Keep victim prone and quiet. EXTERNAL: If skin is contacted, wash immediately with soap and water.

CALL A PHYSICIAN IMMEDIATELY: NOTE TO PHYSICIAN: In case of poisoning, administer barbiturates as for anti-convulsant therapy. Observe patient carefully since repeated treatment may be needed.

(SEE WARNING IN LEFT PANEL)

ACCEPTED

Jun 28 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL

AGRICULTURAL DIVISION



LITTLE ROCK, ARKANSAS

OLIN MATHEWSON CHEMICAL CORPORATION

DIREC

Apply in sufficient quantity
For airplane application
dilute spray solution
growth and good crop

COTTON INSECTS:

Apply when insects are present in sufficient numbers to require necessary or as specified on label—THRIPS and COTTON WEEVIL and COTTON BOLL WORM and 2 to 3 pts. per acre and FALL ARMYWORM on cotton. Workers entering field should wear protective clothing.

WHEAT, OATS, BARLEY

cutworms, Pale weevil, etc. 1 to 2 pints per acre (0.25 lb. coverage. MAKE SURE IT REACHES THE PEAR. Do not apply within 100 yds. of livestock on trees to livestock.

1258-443

ENDRIN

1.6 LB.

EMULSIFIABLE

ACTIVE INGREDIENTS:

Endrin*	19.7%
Methylated aromatic petroleum derivatives	75.4%

INERT INGREDIENTS:

.....	4.9%
Total	100.0%

*Hexachloroepoxyoctahydro-endo, endo-dimethanonaphthalene
Contains 1.6 lbs. Endrin per U. S. gallon at 68° F.

RED

DANGER

KEEP OUT OF REACH OF CHILDREN

POISONOUS BY SKIN CONTACT, INHALATION OR SWALLOWING, RAPIDLY ABSORBED THROUGH SKIN. SPRAY MIST EXTREMELY HAZARDOUS. Do not get in eyes, on skin or on clothing. Do not breathe spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Clean up spillage. Do not use indoors without adequate exhaust ventilation. Use clean clothing daily and wear clean synthetic rubber gloves. In case of contact immediately remove all contaminated clothing and wash skin or eyes with plenty of water for at least 15 minutes; for eyes, get medical attention. Wash clothing before re-use. Avoid contamination of feed or foodstuffs.



ANTIDOTE: INTERNAL: If any of the insecticide is swallowed; give a tablespoon of salt in a glass of warm water. Repeat until vomit fluid is clear. Keep victim prone and quiet. EXTERNAL: If skin is contacted, wash immediately with soap and water.

CALL A PHYSICIAN IMMEDIATELY: NOTE TO PHYSICIAN: In case of poisoning, administer barbiturates as for anti-convulsant therapy. Observe patient carefully since repeated treatment may be needed.

(SEE WARNING IN LEFT PANEL)

DIRECTIONS FOR USE

Apply in sufficient water for good uniform coverage. For airplane applications apply in 3 to 10 gallons dilute spray solution per acre, consistent with crop growth and good coverage.

COTTON INSECTS:

Apply when insects appear in damaging numbers and repeat as necessary or as specified by your State Agricultural Experiment Station—THRIPS and COTTON FLEAHOPPER use 1 pt. per acre; BOLL WEEVIL and COTTON BOLLWORM use 1 pt. per acre early season and 2 to 3 pts. per acre mid-season and late season; LEAFWORMS and FALL ARMYWORM use 1 to 2½ pts. per acre. Do not graze dairy animals or animals being finished for slaughter on treated fields. Workers entering fields within 5 days after treatment should wear protective clothing.

WHEAT, OATS, BARLEY & RYE: For control of Armyworms, Army cutworms, Pale western cutworms and Fall armyworms; apply 1¼ pints per acre (0.25 lb. actual Endrin) in sufficient water for good coverage. MAKE SINGLE APPLICATION WHEN INSECTS FIRST APPEAR. Do not apply within 45 days of harvest. Do not graze livestock on treated forage. Do not feed threshings to livestock.

1258-443

ACCEPTED

Jun 28 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL

tion of law or otherwise,
tion and is reasonably fit
nce with directions under
to the use of this product
s, or in combination with
eable to seller, and buyer
ch use. Seller shall not be
ges arising from a breach

. During Commercial
loading operations,
ask or respirator ap-
or Endrin protection.
als out of operating
drift. Vacated areas
ecticides and volatile

NG
f lakes, ponds, and
on the label. Do not
spray from the crop

ncture and crush be-
containers and dis-
vay from water sup-

LRL94F-06R

GE MINN.
NON PA
TON, TEN
LE ROCK, ARK
IA, NEB

AGRICULTURAL DIVISION
LITTLE ROCK, ARKANSAS



OLIN MATHESSON CHEMICAL CORPORATION

PROCESSED IN ARIZ
INDIANAPOLIS, IND
ST. LOUIS, MO
AUGUSTA, GA
WICHITA, KAN

Right Panel - Endrin 1.6 lb. Emulsifiable

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

WARNING

Do not use or store near heat or open flame. During Commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator approved by the U.S. Department of Agriculture for Endrin protection. Keep unprotected humans and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until the drifting insecticides and volatile residues have dissipated.

PRECAUTIONS IN USING

Highly toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift of spray from the crop area being treated.

DESTROY THIS CONTAINER WHEN EMPTY — Puncture and crush beyond any possible re-use. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies.

USDA Reg. No. 1258-443

LR194F-06R

ENDRIN

1.6 LB.

EMULSIFIABLE

ACTIVE INGREDIENTS:

Endrin* 19.7%
Methylated aromatic petroleum derivatives 75.4%

INERT INGREDIENTS:

Total 100.0%

*Hexachloroepoxyoctahydro-endo, endo-dimethanonaphthalene
Contains 1.6 lbs. Endrin per U. S. gallon at 68° F.

DANGER

KEEP OUT OF REACH OF CHILDREN

POISONOUS BY SKIN CONTACT, INHALATION OR SWALLOWING, RAPIDLY ABSORBED THROUGH SKIN. SPRAY MIST EXTREMELY HAZARDOUS. Do not get in eyes, on skin or on clothing. Do not breathe spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Clean up spillage. Do not use indoors without adequate exhaust ventilation. Use clean clothing daily and wear clean synthetic rubber gloves. In case of contact immediately remove all contaminated clothing and wash skin or eyes with plenty of water for at least 15 minutes; for eyes, get medical attention. Wash clothing before re-use. Avoid contamination of feed or foodstuffs.

POISON

ANTIDOTE: INTERNAL: If any of the insecticide is swallowed; give a tablespoon of salt in a glass of warm water. Repeat until vomit fluid is clear. Keep victim prone and quiet. **EXTERNAL:** If skin is contacted, wash immediately with soap and water.

CALL A PHYSICIAN IMMEDIATELY; NOTE TO PHYSICIAN: In case of poisoning, administer barbiturates as for anti-convulsant therapy. Observe patient carefully since repeated treatment may be needed.

(SEE WARNING IN LEFT PANEL)

DIRECTIONS FOR USE

Apply in sufficient water for good uniform coverage. For airplane applications apply in 3 to 10 gallons dilute spray solution per acre, consistent with crop growth and good coverage.

COTTON INSECTS:

Apply when insects appear in damaging numbers and repeat as necessary or as specified by your State Agricultural Experiment Station—**THRIPS** and **COTTON FLEAHOPPER** use 1 pt. per acre; **BOLL WEEVIL** and **COTTON BOLLWORM** use 1 pt. per acre early season and 2 to 3 pts. per acre mid-season and late season; **LEAFWORMS** and **FALL ARMYWORM** use 1 to 2½ pts. per acre. Do not graze dairy animals or animals being finished for slaughter on treated fields. Workers entering fields within 5 days after treatment should wear protective clothing.

WHEAT, OATS, BARLEY & RYE: For control of Armyworms, Army cutworms, Pale western cutworms and Fall armyworms; apply 1¼ pints per acre (0.25 lb. actual Endrin) in sufficient water for good coverage. **MAKE SINGLE APPLICATION WHEN INSECTS FIRST APPEAR.** Do not apply within 45 days of harvest. **Do not graze livestock on treated forage. Do not feed threshings to livestock.**

1258-443

ACCEPTED
Jun 28 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PURPOSES

SAVAGE MINN.
LEBANON, PA
HOUSTON, TEX
NO. LITTLE ROCK, ARK
OMAHA, NEB

AGRICULTURAL DIVISION **Olin**
LITTLE ROCK, ARKANSAS

OLIN MATTHEWSON CHEMICAL CORPORATION

DIRECTOR, ARK.
LITTLE ROCK, ARK.
SAC, HOUSTON, TEX.
SAC, OMAHA, NEB.
SAC, WICHITA, KAN.

Right Panel - Endrin 1.6 lb. Emulsifiable

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions, not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

ENDRIN

1.6 LB.

EMULSIFIABLE

DIRECTIONS FOR USE

Apply in sufficient water for good uniform coverage. For airplane applications apply in 3 to 10 gallons dilute spray solution per acre, consistent with crop growth and good coverage.

WARNING

Do not use or store near heat or open flame. During Commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator approved by the U.S. Department of Agriculture for Endrin protection. Keep unprotected humans and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until the drifting insecticides and volatile residues have dissipated.

ACTIVE INGREDIENTS:

Endrin*	19.7%
Methylated aromatic petroleum derivatives	75.4%
INERT INGREDIENTS:	
Total	4.9%
	100.0%

*Hexachloroepoxyoctahydro-endo, endo-dimethanonaphthalene
Contains 1.6 lbs. Endrin per U. S. gallon at 68° F.

COTTON INSECTS:

Apply when insects appear in damaging numbers and repeat as necessary or as specified by your State Agricultural Experiment Station--THRIPS and COTTON FLEAHOPPER use 1 pt. per acre; BOLL WEEVIL and COTTON BOLLWORM use 1 pt. per acre early season and 2 to 3 pts. per acre mid-season and late season; LEAFWORMS and FALL ARMYWORM use 1 to 2½ pts. per acre. Do not graze dairy animals or animals being finished for slaughter on treated fields. Workers entering fields within 5 days after treatment should wear protective clothing.

PRECAUTIONS IN USING

Highly toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift of spray from the crop area being treated.

DESTROY THIS CONTAINER WHEN EMPTY -- Puncture and crush beyond any possible re-use. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies.

USDA Reg. No. 1258-443

LRI94F-06R

DANGER
KEEP OUT OF REACH OF CHILDREN

POISONOUS BY SKIN CONTACT, INHALATION OR SWALLOWING, RAPIDLY ABSORBED THROUGH SKIN. SPRAY MIST EXTREMELY HAZARDOUS. Do not get in eyes, on skin or on clothing. Do not breathe spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Clean up spillage. Do not use indoors without adequate exhaust ventilation. Use clean clothing daily and wear clean synthetic rubber gloves. In case of contact immediately remove all contaminated clothing and wash skin or eyes with plenty of water for at least 15 minutes; for eyes, get medical attention. Wash clothing before re-use. **Avoid contamination of feed or foodstuffs.**




ANTIDOTE: INTERNAL: If any of the insecticide is swallowed, give a tablespoon of salt in a glass of warm water. Repeat until vomit brings it out. Keep victim prone and quiet. **EXTERNAL:** If skin is contacted, wash immediately with soap and water. **CALL A PHYSICIAN IMMEDIATELY. NOTE TO PHYSICIANS:** In case of poisoning, administer barbiturates as for anti-convulsant therapy. Observe patient carefully since repeated treatment may be needed.

(SEE WARNING IN LEFT PANEL)

WHEAT, OATS, BARLEY & RYE: For control of Armyworms, Army cutworms, Pale western cutworms and Fall armyworms; apply 1½ pints per acre (0.25 lb. actual Endrin) in sufficient water for good coverage. **MAKE SINGLE APPLICATION WHEN INSECTS FIRST APPEAR.** Do not apply within 45 days of harvest. **Do not graze livestock on treated forage. Do not feed threshings to livestock.**

1258-443



Right Panel - Endrin 1.6 lb. emulsifiable