



CONTENTS 4 LBS. NET

MALATHION 5% DUST

FOR USE ON PETS, ORNAMENTALS AND VEGETABLES

ACTIVE INGREDIENTS:

Malathion* 5.0%

INERT INGREDIENTS: 95.0%

Total 100.0%

*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not contaminate feed, utensils or food-stuffs. Avoid prolonged breathing of the dust. Avoid prolonged or repeated contact with skin; wash thoroughly after using.

NOTE TO PHYSICIAN: Malathion, upon repeated, prolonged careless use may cause cholinesterase inhibition. Atropine is antidotal.

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.



May 1, 1969

FEDERAL INSECTICIDE
REGISTRATION ACT
REGISTERED
1258-468

MALATHION 5% DUST

DIRECTIONS FOR USE

For best results use a good garden type or puffer type duster, using enough dust to lightly but thoroughly cover the foliage.

BEANS, PEAS, GRAPES, STRAWBERRIES, CABBAGE, BROCCOLI, KALE, MUSTARD, SPINACH, TURNIPS, LETTUCE, ONIONS and POTATOES: For the control of aphids, spider mites, Mexican bean beetle, imported cabbage worm, cabbage looper, lace bug, onion thrip, and potato leafhopper — apply with thorough coverage and repeat as necessary. CAUTION: Do not use on leaf lettuce within 14 days of harvest or within 7 days of harvest of head lettuce or other crops listed.

ORNAMENTALS: For the control of aphids, mealy bugs, spider mites, white fly, tarnished plant bug, thrip, rose leafhopper, boxwood miner, bagworm, azalea scale, pine leaf scale, and Japanese beetles; apply with thorough coverage and repeat as necessary. Avoid use on certain ferns including Boston, Maidenhair and Pteris as well as on some species of Crassula.

DAIRY CATTLE: For the control of horn flies, apply 3 tablespoonfuls of 5% dust per animal. Apply the dust on the back and neck of the animals and repeat applications at 10 to 14 day intervals. Do not apply to dairy cattle later than five hours before milking or during milking. Do not contaminate feed, food containers and water troughs.

FLIES: As an aid in the reduction of flies, dust around trash cans, garbage cans and breeding areas as needed. Breeding sites such as manure or other waste material should be eliminated. Do not use in dwellings or in milk rooms.

ANIMAL PESTS: For control of fleas on dogs and cats use enough dust to cover the animal lightly but thoroughly, then work into the hair coat using a soft-bristle brush. The neck hair should receive slightly heavier application of dust. If animals run free, reinfestations may occur and retreatments may be necessary every two or three weeks. Examine pets closely for reinfestations to avoid unnecessary treatments.

As an aid in the reduction of fleas in dog beds, kennels, pens and yards — first clean the area then dust thoroughly using 1 to 2 pounds per 1000 square feet. Pay special attention to areas where insects may be hiding.

USE PRECAUTIONS

This material is toxic to fish. Keep out of lakes, ponds and streams. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Extension Service. Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift of spray from the crop area being treated.

Destroy empty bags by burning (KEEP OUT OF SMOKE).

USDA Reg. No. 1258-468

LR153E-03S



ACCEPTED WITH COMMENTS



CONTENTS 4 LBS. NET

FOR USE ON PETS, ORNAMENTALS AND VEGETABLES

ACTIVE INGREDIENTS:

Malathion* 5.0%

INERT INGREDIENTS:

95.0%

Total 100.0%

O, O dimethyl dithiophosphate of diethyl mercaptosuccinate

NOTICE TO BUYER

DIRECTIONS FOR USE

BEANS, PEAS, GRAPES, STRAWBERRIES, CABBAGE, BROCCOLI, KALE, MUSTARD SPINACH, TURNIPS, LETTUCE, ONIONS and POTATOES:

Apply this material to the foliage of the plants at the rate of 1/2 to 1 quart per acre. Repeat applications at 7 to 10 day intervals. Do not apply to plants in bloom. Do not apply to plants in fruit. Do not apply to plants in seed. Do not apply to plants in storage. Do not apply to plants in transit. Do not apply to plants in processing. Do not apply to plants in distribution. Do not apply to plants in retail.

ORNAMENTALS: Apply this material to the foliage of the plants at the rate of 1/2 to 1 quart per acre. Repeat applications at 7 to 10 day intervals. Do not apply to plants in bloom. Do not apply to plants in fruit. Do not apply to plants in seed. Do not apply to plants in storage. Do not apply to plants in transit. Do not apply to plants in processing. Do not apply to plants in distribution. Do not apply to plants in retail.

DAIRY CATTLE: For the control of horn flies, apply this material at the rate of 1/2 to 1 quart per animal. Apply the material to the animal's body at the rate of 1/2 to 1 quart per application. Repeat applications at 7 to 10 day intervals. Do not apply to animals in bloom. Do not apply to animals in fruit. Do not apply to animals in seed. Do not apply to animals in storage. Do not apply to animals in transit. Do not apply to animals in processing. Do not apply to animals in distribution. Do not apply to animals in retail.

FLEEC: Apply this material to the fleece of the animals at the rate of 1/2 to 1 quart per animal. Apply the material to the animal's body at the rate of 1/2 to 1 quart per application. Repeat applications at 7 to 10 day intervals. Do not apply to animals in bloom. Do not apply to animals in fruit. Do not apply to animals in seed. Do not apply to animals in storage. Do not apply to animals in transit. Do not apply to animals in processing. Do not apply to animals in distribution. Do not apply to animals in retail.

ANIMAL PESTS: Apply this material to the animal's body at the rate of 1/2 to 1 quart per animal. Apply the material to the animal's body at the rate of 1/2 to 1 quart per application. Repeat applications at 7 to 10 day intervals. Do not apply to animals in bloom. Do not apply to animals in fruit. Do not apply to animals in seed. Do not apply to animals in storage. Do not apply to animals in transit. Do not apply to animals in processing. Do not apply to animals in distribution. Do not apply to animals in retail.

Apply this material to the animal's body at the rate of 1/2 to 1 quart per animal. Apply the material to the animal's body at the rate of 1/2 to 1 quart per application. Repeat applications at 7 to 10 day intervals. Do not apply to animals in bloom. Do not apply to animals in fruit. Do not apply to animals in seed. Do not apply to animals in storage. Do not apply to animals in transit. Do not apply to animals in processing. Do not apply to animals in distribution. Do not apply to animals in retail.

USE PRECAUTIONS

This material is toxic to fish. Keep out of lakes, ponds and streams. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Extension Service. Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift of spray from the crop area being treated.

Destroy empty bags by burning. KEEP OUT OF SMOKE

AGRICULTURAL DIVISION **Olin**

LITTLE ROCK, ARKANSAS
OLIN MATHIESON CHEMICAL CORPORATION
LEBANON, PA. OMAHA, NEB. ST. LOUIS, MO.
HOUSTON, TEX. PHOENIX, ARIZ. AUGUSTA, GA.
NO. LITTLE ROCK, ARK. RICHMOND, IND. WICHITA, KAN.
SAVAGE, MINN.

AGRICULTURAL DIVISION **Olin**

LITTLE ROCK, ARKANSAS
OLIN MATHIESON CHEMICAL CORPORATION