




TERRACLO[®]

SOIL FUNGICIDE

10% GRANULAR

Controls Certain Soil-borne Diseases of Lawns and Cotton



ACTIVE INGREDIENT:

Pentachloronitrobenzene 10%

INERT INGREDIENTS: 90%

Total 100%

TERRACLO[®] is a trademark of Olin Mathieson Chemical Corporation.

CAUTION

Avoid prolonged inhalation of dust. This material may cause skin irritation. Wash hands and face thoroughly after using. Keep out of the reach of children. Avoid contamination of feed and foodstuffs.

DIRECTIONS FOR USE

Lawns (Brown Patch—Rhizoctonia solani): For Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses: Make one application in the fall or spring at first indication of infection. Apply 7½ lbs. per 1000 sq. ft. Following application, the treated area should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded,

or if the disease is severe or reappears, the area should be retreated 3-4 weeks later. Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

For Northern or cool season lawns such as Rye Grass, Fescue, Bluegrass, and similar grasses: Apply 2 to 2.5 lbs. at first sign of disease. Repeat every 7-10 days during warm, damp weather. Water treated area lightly to wash chemical down to soil.

Cotton (Damping-off, Seed and Seedling Rots—Rhizoctonia): Furrow Application: Apply 10-20 lbs. per acre to the planting furrow. Apply over seed and covering soil at planting time, using suitable application equipment.

IMPORTANT NOTICE

Since the use of this product is beyond the control of the seller, seller makes no warranty of any kind, express or implied, concerning the use of this product. User assumes all risk of use or handling whether in accordance with directions or not.

NET CONTENTS LBS.

CHEMICALS DIVISION 
The Developers of PCNB Fungicides

1258-555

07/12/1962

111