

Terraclor® 10% GRANULAR

SOIL FUNGICIDE

Controls Certain Soil-Borne Diseases Of
BEANS • COTTON • CRUCIFERS
PEANUTS • POTATOES
LAWNS • ORNAMENTALS

NOTICE TO BUYER

Seller warrants the warranty, express or implied, by operation of law, or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special or consequential or incidental damages arising from a breach of this warranty.

USE CAUTIONS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil.

Dispose of empty bags by burning. (KEEP OUT OF SMOKE).

DIRECTIONS FOR USE

BEANS (Root and Stem Rot — *Rhizoctonia solani*):

PRE-TREATMENT: Apply 10 to 15 lbs. per acre 14 days before planting. **POST-TREATMENT:** Apply 8 to 10 lbs. per acre 14 days before planting. **NOTE:** Do not feed treated vegetable stock.

COTTON (Damping-off, Seed and Seedling Rots — *Rhizoctonia solani*):

For furrow and broadcast apply 10 to 15 lbs. per acre 14 days before planting. For 2 lbs. per 1000 seedlings in a field 40 rows wide. Apply over seed and cover with soil and plant the seed. Use suitable application equipment for furrow with the seed and broadcast to all the vegetable stock.

CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, AND BROCCOLI (Clubroot — *Plasmodiophora brassicae*):

Apply 200 lbs. per 100 feet of row for broadcast and Brussels sprouts, cauliflower and cabbage. Apply 100 lbs. per 100 feet of row for cauliflower and Brussels sprouts. Apply 100 lbs. per 100 feet of row for cauliflower and Brussels sprouts. Apply 100 lbs. per 100 feet of row for cauliflower and Brussels sprouts. Apply 100 lbs. per 100 feet of row for cauliflower and Brussels sprouts.

CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, AND BROCCOLI (Clubroot — *Plasmodiophora brassicae*):

BROADCAST: Apply 200 lbs. per acre to row.

ACTIVE INGREDIENTS:

Pentachloronitrobenzene	10%
INERT INGREDIENTS:	90%
Total	100%

Terraclor® is a trademark of Olin Corporation.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhaling dust. This material may cause skin irritation. Wash hands and face thoroughly after using. Wear gloves and a respirator when loading the spreader or duster.

Do not contaminate feed or foodstuffs.

In transplanting, dig and press dust into bottom of hole. **NOTE:** For control of cabbage, cauliflower, and Brussels sprouts, it is very important to thoroughly mix with soil.

CAULIFLOWER (Wire Stem or Black root — *Corticium solani*):

Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row.

PEANUTS (Southern blight — *Sclerotium rolfsii*):

Treatment: Apply 10 lbs. per 1000 feet of row. Apply 10 lbs. per 1000 feet of row. Apply 10 lbs. per 1000 feet of row. Apply 10 lbs. per 1000 feet of row. Apply 10 lbs. per 1000 feet of row. Apply 10 lbs. per 1000 feet of row.

POTATOES (Rhizoctonia Solani):

Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row.

POTATOES (Rhizoctonia solani):

Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row.

POTATOES (Rhizoctonia solani):

Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row.

POTATOES (Rhizoctonia solani):

Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row. Apply 100 lbs. per 1000 feet of row.

For the tubers and roots, the soil should be treated to make treatment effective.

LAWNS (Brown patch — *Rhizoctonia solani*):

For warm season lawns (St. Augustine, Bermuda and similar grasses) make one application in the fall and one in the spring. Apply 7 lbs. per 1000 sq. ft. Following application, the treated area should be kept watered to insure movement through the grass to soil level. If the treated area is subjected to unusual heavy rainfall or topped, or if the disease is severe or reappears, the area should be retreated 3 to 4 weeks later. For cool season lawns (such as Ryegrass, Fescue, Bluegrass, Bent and similar grasses) apply 2 to 2.5 lbs. per 1000 sq. ft. at first sign of disease. Repeat every 7 to 10 days during warm and wet weather. Water treated area daily to wash off dust. Note: A permanent growing of this is a hindrance to the application of the dust. Grass may be cut 1/2 inch high and will disappear in short time.

ORNAMENTALS

AFRICAN VIOLETS, BEGONIAS, CARNATIONS, CHRYSANTHEMUMS, POINSETTIAS, SNAPDRAGONS, and similar bedding plants (Stem Rot — *Rhizoctonia*):

Apply 10 lbs. per 1000 sq. ft. of area. Apply 10 lbs. per 1000 sq. ft. of area. Apply 10 lbs. per 1000 sq. ft. of area. Apply 10 lbs. per 1000 sq. ft. of area. Apply 10 lbs. per 1000 sq. ft. of area. Apply 10 lbs. per 1000 sq. ft. of area.

BULBOUS IRIS, HYACINTH, NARCISSUS, AND TULIP (Crown Rot and Black Rot):

Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area.

CALENDULA, LARKSPUR, SNAPDRAGON, AND SWEET PEA (Stem Rot — *Sclerotinia*):

Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area.

DUTCH IRIS (Bulb and stem rot caused by *Sclerotium rolfsii*):

Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area.

EASTER LILIES (Rhizoctonia root rot):

Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area.

GLADIOLUS (Neck or Bulb Dry Rot — *Stromatinia* or *Sclerotinia gladioli*):

Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area. Apply 100 lbs. per 1000 sq. ft. of area.

LR392F-12U

EPA Reg. No. 1258-555-AA

Accepted for sale
 Sept 6, 1970