GENERAL INFORMATION

If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using.

When filling spray tank first fill tank ¼ full of water; add the calculated amount of 2,4-D emulsifiable, then finish filling tank with water with by-pass open and pump running.

Material may tend to settle when not agitated. When mixture is left in sprayer allow by-pass pressure to re-agitate before resuming spraying.

PRECAUTION

BECAUSE OF THE DIFFICULTIES OF REMOVING 2,4-D FROM SPRAYERS, DO NOT USE THE SAME SPRAY EQUIPMENT FOR APPLYING OTHER MATERIALS TO PLANTS AS INJURY WILL RESULT. USE A SEPARATE SPRAYER FOR THIS PURPOSE.

Do not reuse empty container. Destroy by burying in a safe place. Do not burn.

LR319G-01W 7324711
EPA Reg. No. 1258-652

SALVO

(LOW VOLATILE)

WEED KILLER

ACTIVE INGREDIENT:	
2-ethylhexyl ester of 2,4-Dichlorophenoxyacetic Acid	**85.9%**
INERT INGREDIENTS:	**14.1%**
Total	**100.0%**
Equivalent to 2,4-Dichlorophenoxyacetic Acid	**57.0%**

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION. Avoid contact with eyes, skin and clothing. Do not take internally. Do not contaminate feed or foodstuffs.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc (Course sprays are less likely to drift.)

Do not contaminate domestic or irrigating water supplies.

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not seed lawns gardens or fields until at least four weeks after application of spray to area. Do not mix with liquid fertilizers. Do not apply in any manner not specified on the label.

DO NOT STORE NEAR FERTILIZERS, SEEDS, INSECTICIDES, FUNGICIDES OR PLANTS

TIME OF APPLICATION

Annual weeds are most susceptible in the seedling stage and early growth stages of development and when conditions are such as to promote a vigorous growth.

WHEN USED IN CROP SPRAYING AS A SELECTIVE SPRAY, THE STAGE OF GROWTH OF THE CROP MUST ALSO BE CONSIDERED, OTHERWISE SEVERE DAMAGE MAY RESULT

SOME SUSCEPTIBLE WEEDS

Arrowhead	Henbit	Prickly
Artichoke	Horsetail	P
Beggar Ticks	Jimsonweed	
Bull Thistle	Kochia	
Butterprint	Lambs	
Catnip		
Chickory		(annual)
Cocklebur		seed
Dock		Tomacs
		Sunflower
		Wild Mustard
		Wild Onion
	Pepper Grass	Wild Parsnip
	Pig Weed	Wild Radish
Wild Rape		Wild Sweet Potato

This section to be amended as per sheet attached

Some weeds above, listed as susceptible may not be completely killed but growth is so retarded by proper application that they are no longer a factor affecting growth and yield of the crop.

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damage arising from a breach of this warranty.

CROP	*Amount SALVO Per Acre	APPLICATION	CAUTION
CORN Pre-emergence	1-1 2 2-1 4 pints	Spray uniformly over the soil surface after planting (preferably 3 to 5 days) but before emergence. Use in 5 to 20 gallons of water.	Do not use on light sandy soils, or when moisture is inadequate for normal weed growth.
CORN, MILO Postemergence	1 5 2 5 pt.	Apply when crop is small and more resistant to 2,4-D but when weeds are succulent or in active growth stage. Use lower dosage only against succulent, non-resistant annual broadleaf weeds.	Use drop nozzles when crop is 10 inches tall or higher. Keep spray off leaves. Do not spray after tassels appear. Some injury to corn may occur.
WHEAT OATS BARLEY RYE	FALL 2 5 4 5 pt. SPRING 2 5 1 2 pt.	In spring when weeds are small and grain is in full tiller stage (4 to 8 inches high) and always before boot stage. After plants are fully tillered but before stems begin to joint	Do not apply during boot or dough stage. Do not forage or graze treated grain fields within two weeks after treatment. Do not feed treated straw to livestock
PERENNIAL GRASS PASTURES, MEADOWS	4 5 pt.	In early spring when sufficient weeds have emerged and are actively growing and not too mature	Injury may result to bent grass and to clovers. Usually clovers will recover from early spring applications. Do not apply when grass is in boot to milk stage. Do not apply to newly seeded areas or after heading begins. Do not graze treated areas to dairy animals within 7 days after treatment.
LAWNS, ORNAMENTAL TURF GRASS & GOLF FAIRWAYS	Large areas 1-1 2 pints per acre. Small areas 3-1 3 teaspoonsful in 3 to 5 gallons of water applied to 1000 square feet.		Spray 3 to 5 days after mowing. Retreatment of perennial weed areas may be necessary. May cause injury to bent grass, carpet grass, dichondra, St. Augustine and to clovers. Protect ornamentals from spray drift

*Application rates may vary slightly from State to State depending upon soil types and weather factors. Consult your State agricultural experiment station for recommendations under local conditions or more specific uses.

SALVO is a trademark of Olin Corporation

AGRICULTURAL DIVISION Olin
LITTLE ROCK ARKANSAS
OLIN CORPORATION