

SALVO®

(LOW VOLATILE) WEED KILLER

ACTIVE INGREDIENT:

Isooctyl ester of 2,4-Dichlorophenoxyacetic Acid† 81.8%

INERT INGREDIENTS: 18.2%

TOTAL **100.0%**

Equivalent to 2,4-Dichlorophenoxyacetic Acid 54.2%

† Isomer Specific by A.O.A.C. Method 6 D01 †

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION. Avoid contact with eyes, skin and clothing. Do not take internally. Do not contaminate feed or foodstuffs.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Course sprays are less likely to drift.) Do not contaminate domestic or irrigating water supplies. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not seed lawns, gardens or fields until at least four weeks after application of spray to area. Do not mix with liquid fertilizers. Do not apply in any manner not specified on the label.

TIME OF APPLICATION

Annual weeds are most susceptible in the seeding stage and early growth stages of development and when conditions are such as to promote a vigorous growth. **WHEN USED IN CROP SPRAYING AS A SELECTIVE SPRAY THE STAGE OF GROWTH OF THE CROP MUST ALSO BE CONSIDERED. OTHERWISE SEVERE DAMAGE MAY RESULT.**

SOME SUSCEPTIBLE WEEDS

Annual Weeds	King Dogbane	Puncture Vine
Antelope	Lotus	Russet
Broomrape	Parrot Pea	Ragweed
Buckwheat	Parrot Pea	Russet Thistle
Blue Thistle	Parrot Pea	Shepherd's Purse
Buttercup	Parrot Pea	Smart Weed
Canada Thistle	Parrot Pea	Sow Thistle (annual)
Cat's Paw	Parrot Pea	Stinkweed
Chickweed	Parrot Pea	Sumac
Chenopod	Parrot Pea	Sunflower
Cocklebur	Parrot Pea	Wild Garlic
Dandelion	Parrot Pea	Wild Mustard
Drumstick	Parrot Pea	Wild Onion
Fern	Parrot Pea	Wild Parsnip
Field Bindweed	Parrot Pea	Wild Radish
Flourish	Parrot Pea	Wild Rape
Groundnut	Parrot Pea	Wild Sweet Potato
Groundsel	Parrot Pea	Yellow Goatbeard

Some weeds above, listed as susceptible may not be completely killed but growth is so retarded by proper application that they are no longer a factor affecting growth and yield of the crop.

NOTICE TO BUYER

This product is a selective herbicide and is for the use in lawns, gardens, etc. as indicated on the label. Do not use on crops or other plants not listed on the label. Do not use on lawns, etc. if the grass is not in the seedling stage. Do not use on lawns, etc. if the grass is not in the seedling stage. Do not use on lawns, etc. if the grass is not in the seedling stage.

LR652-02E

APPLICATION	CAUTION
Apply uniformly over the soil surface after planting (preferably 4 to 5 days) but in emergency use in 5 to 20 gallons of water.	Do not use on light sandy soils or when moisture is inadequate for normal weed growth.
When crop is small and more resistant to 2,4-D but when weeds are in an active growth stage. Use lower dosage only against weeds that are not hard to control.	Use drop nozzles when crop is in the early stages of growth. Keep spray off leaves. Do not spray after 10:00 a.m. Some injury to corn may occur.
Apply when weeds are small and grass is in the seedling stage. Always be sure to use the correct dosage.	Do not apply during boot or dough stage. Do not graze or mow treated areas for 2 weeks after treatment. Do not feed treated straw to livestock.
Apply when weeds are small and grass is in the seedling stage. Always be sure to use the correct dosage.	Injury may result to corn grass and to clover. Usually, injury will be minimal if early spring applications. Do not apply when grass is in boot to milk stage. Do not apply to newly seeded areas. After the seedling stage, do not graze treated areas for 2 weeks after treatment.
Apply when weeds are small and grass is in the seedling stage. Always be sure to use the correct dosage.	Do not use on lawns, etc. if the grass is not in the seedling stage. Do not use on lawns, etc. if the grass is not in the seedling stage.

