

SEVIN* 10% DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 10%

INERT INGREDIENTS:

90%

TOTAL

100%

* Trademark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. DO NOT BREATHE DUST. DO NOT TAKE INTERNALLY. SKIN CONTACT MAY BE HARMFUL. WASH OFF IMMEDIATELY. CONTACT WITH SKIN: WASH HANDS BEFORE EATING. WASH FACE AND HAIR AFTER HANDLING. WEAR REGULAR LONG SLEEVED CLOTHING AND GLOVES. CLEAN CLOTHING DAILY. AVOID CONTAMINATION OF FOOD AND FEEDS.

For more information on the use of Sevin 10% Dust, contact your local distributor or write to Union Carbide Corporation, P.O. Box 117, Little Rock, Arkansas 72202.

USE PRECAUTIONS

Destroy empty bags by burning (KEEP OUT OF SMOKE).

For protection of honey bees avoid use, if possible, during periods when bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms of the label and subject thereto. The Buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

DIRECTIONS FOR USE

TOBACCO (FIELD): For control of tobacco hornworm, budworm, fleabeetles, June beetles and Japanese beetles apply 10 to 15 lbs. per acre at the first sign of infestation and repeat as needed. When dusting for budworm control direct some of the dust into each bud. Do not apply within 3 days of priming or cutting. If late applications are necessary, treat immediately after priming, rather than before priming.

NOTE: Some damage may occur to tender foliage in the presence of rain or high humidity of several days duration. Avoid contacting plant juices.

DIRECTIONS FOR USE (Cont.)

ALFALFA, CLOVER, PEANUTS, SOYBEANS, COWPEAS: For control of blister beetle and Mexican bean beetle use 5 to 10 lbs. per acre. For control of thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetle, bean leaf beetle, Japanese beetle, alfalfa caterpillar, velvetbean caterpillar and green cloverworm use 10 lbs. per acre. For control of armyworms, webworms, corn earworm, grasshoppers and stink bug use 10 to 14 lbs. per acre. (Some leaf injury may occur if applications are made when tender foliage is wet.)

RICE, FORAGE GRASS and PASTURE: For control of thrips, armyworms, stink bugs and grasshoppers use 10 to 14 lbs. per acre. Do not apply within 14 days of harvest of rice.

COTTON: For early-season control of cotton fleahopper, thrips and flea beetles use 5 to 10 lbs. per acre. For MID and LATE season control of boll weevil, bollworm, fall armyworm, leaf perforator, leaf rollers, grasshoppers, leafhoppers and tarnished plant bug use 10 to 20 lbs. per acre. For salt marsh caterpillar use 20 lbs. per acre. Allow 7 days between last application and grazing treated fields or harvesting crop residues for use as feed or bedding for dairy or meat animals.

EGGPLANT, PEPPERS, TOMATOES: For control of flea beetles and leafhoppers use 5 to 10 lbs. per acre. For control of tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bugs, lacebugs and lygus bugs use 10 to 20 lbs. per acre.

CUCUMBERS, SQUASH, PUMPKINS, MELONS: For control of pickleworm and melonworm use 6 to 9 lbs. per acre. Use the higher rate for control of cucumber beetles, squash bug, flea beetles and leafhoppers.

STRAWBERRIES: For control of meadow spittlebug, leaf roller and strawberry weevil use 10 to 18 lbs. per acre. Do not apply within 24 hours of harvest.

BEANS: For control of Mexican bean beetle, leafhoppers, bean leaf beetle, cucumber beetle, flea beetles, Japanese beetle and velvetbean caterpillar use 10 lbs. per acre. For corn earworm, armyworms, stink bugs and tarnished plant bugs use 10 to 15 lbs. per acre. When label directions and limitations are followed, forage, vines and hay may be fed to meat and dairy animals.

LR338E-05Q 602349
USDA Reg. No. 1258-692

SAVAGE, MINN.
BALTIMORE, MD.
HOUSTON, TEX.
NO. LITTLE ROCK, ARK.
OMAHA, NEB.

AGRICULTURAL DIVISION **Olin**
LITTLE ROCK, ARKANSAS

OLIN MATHIESON CHEMICAL CORPORATION

PHOENIX, ARIZ.
RICHMOND, IND.
ST. LOUIS, MO.
WILLIAMSTON, N.C.
WICHITA, KAN.

SEVIN 10 DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 10%

INERT INGREDIENTS:

90%

TOTAL

100%

Sevin 10 Dust contains 10% Carbaryl (1-naphthyl N-methylcarbamate) and 90% inert ingredients.

USE PRECAUTIONS

Destroy empty bags by burning (KEEP OUT OF SMOKE).

For protection of honey bees avoid use, if possible, during periods when bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

NOTICE TO BUYER

This product is to be limited to the terms of the label. It is not to be used on any crop, animal, or property not specified on the label. It is not to be used in any manner not specified on the label.

TOBACCO (FIELD): For control of tobacco rootworm, apply 10 lbs. per acre at the first sign of infestation and repeat as needed. For control of tobacco rootworm, apply 10 lbs. per acre at the first sign of infestation and repeat as needed. For control of tobacco rootworm, apply 10 lbs. per acre at the first sign of infestation and repeat as needed.

NOTE: Do not apply to any crop or animal not specified on the label. Do not apply to any crop or animal not specified on the label.

ALFALFA, CLOVER, PEANUTS, SOYBEANS, COWPEAS: For control of alfalfa beetle and Mexican bean beetle use 5 to 10 lbs. per acre. For control of thrips, leafhoppers, tree-toad, ground sallow, hopper, locust, green stink bug, leaf beetle, Japanese beetle, alfalfa caterpillar, velvetbean caterpillar, and green cloverworm use 10 lbs. per acre. For control of armyworms, cutworms, corn earworm, grasshoppers, and ticks use 10 to 14 lbs. per acre. Do not apply to any crop or animal not specified on the label.

RICE, FORAGE GRASS and PASTURE: For control of rice thrips, grasshoppers, and grasshoppers use 10 to 14 lbs. per acre. Do not apply within 10 days of harvest.

COTTON: For control of cotton rootworm, cotton leafhopper, cotton bollworm, and cotton bollworm use 10 lbs. per acre. For control of all other cotton pests use 10 to 14 lbs. per acre. Do not apply to any crop or animal not specified on the label.

EGGPLANT, PEPPERS, TOMATOES: For control of eggplant and pepper thrips use 5 to 10 lbs. per acre. For control of all other eggplant and pepper pests use 10 to 14 lbs. per acre. Do not apply to any crop or animal not specified on the label.

CUCUMBERS, SQUASH, PUMPKINS, MELONS: For control of cucumber and squash beetle use 10 lbs. per acre. For control of all other cucumber, squash, pumpkin, and melon pests use 10 to 14 lbs. per acre.

STRAWBERRIES: For control of strawberry rootworm use 10 lbs. per acre. For control of all other strawberry pests use 10 to 14 lbs. per acre.

BEANS: For control of Mexican bean beetle use 5 to 10 lbs. per acre. For control of all other bean pests use 10 to 14 lbs. per acre. Do not apply to any crop or animal not specified on the label.

U.S. Pat. 2,616,644
A. Reg. No. 125,644

