



## MIXING INSTRUCTIONS

"ANSAR" 170 Herbicide must be thoroughly dissolved. Fill the tank about half full with water, and with the pump operating, add the required amount of "ANSAR" 170 Herbicide. Finish filling the tank with water.

If preferred, "ANSAR" 170 Herbicide can be premixed by dissolving the required amount of "ANSAR" 170 Herbicide and suitable surfactant in 3 to 5 gallons of water in an open head bucket. This premix should then be placed in the partially filled spray tank, and the tank filled.

## DIRECTIONS FOR USE

### GENERAL GRASS AND WEED CONTROL IN NON-CROP AREAS

"ANSAR" 170 Herbicide is useful for controlling Johnson Grass, nut grass, water grass, goose grass and other weeds on drainage ditch banks, right-of-ways, implement storage yards, fence rows and similar non-crop areas.

Mix "ANSAR" 170 Herbicide at a rate of 3 pints, plus 1½ to 2 quarts of Olin Surfactant 80 in 50 gallons of water. Spray the unwanted vegetation thoroughly to the point of run-off. If re-growth occurs, retreat as required. KEEP LIVESTOCK OFF TREATED AREAS.

ANSAR® is a registered trademark of The Ansul Company, Marinette, Wisconsin

# ANSAR® 170

## HERBICIDE

### FOR POST-EMERGENCE WEED CONTROL IN COTTON And IN NON-CROP AREAS

ACTIVE INGREDIENT:	
Monosodium Acid Methanearsonate	50.6%
INERT INGREDIENTS	49.4%
Total arsenic (as elemental) all in water soluble form	23.41%
(Contains the equivalent of 6.6 lbs. MSMA per gallon at 68° F.)	

(12 pt) **CAUTION** (18 pt) **KEEP OUT OF REACH OF CHILDREN** RED

Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin, wash thoroughly after using. Avoid contamination of feed and foodstuffs.

DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION OR BY ANIMALS, WILDLIFE AND AQUATIC LIFE, OR WATER USED FOR IRRIGATION PURPOSES.

Do not use this product in any manner not specified on this label, since crop residues and injury may result.

Do not store spray solution in tank for a prolonged period, and clean application equipment thoroughly after use by flushing with water.

## DIRECTIONS FOR USE (Con't.)

**COTTON** "ANSAR" 170 Herbicide is useful as a directed spray application to weeds in cotton for post-emergence control of weeds and grasses including Johnson grass, nut grass, water grass, goose grass, dallis grass and cocklebur. Mix at a rate of 3 pints "ANSAR" 170 Herbicide, plus 1½ to 2 quarts of Olin Surfactant 80, in 50 gallons of water. Apply at a broadcast rate of 40 gallons of solution per acre. For band application, apply at the rate of one gallon of solution for each 1 inch of band width to be treated, on cotton grown in 40 inch rows. A second application may be required during humid growing conditions. Time the second application, if necessary, about one to three weeks after the first application. APPLY ONLY WHEN COTTON IS 3 INCHES HIGH TO FIRST BLOOM. DO NOT APPLY AFTER FIRST BLOOM. DO NOT FEED TREATED FOLIAGE TO LIVESTOCK, OR GRAZE TREATED AREAS.

Applications should be made with a low volume sprayer having satisfactory pumping and bypass action. Nozzles should be placed so as to avoid spraying the cotton but also giving good coverage of the weeds and grasses. PROPER COVERAGE IS VERY IMPORTANT. Adjust nozzles to allow maximum coverage of weeds and grasses with minimum contact to the cotton plant. KEEP SPRAY OFF COTTON FOLIAGE.

Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment but the plant will develop normally.

**IMPORTANT NOTICE:** Since the use of this product is beyond the control of the seller, seller makes no warranty, express or implied, concerning the use of this product. User assumes all risk of use or handling, whether in accordance with directions or not.

LR359B-01Q 6224336  
USDA Reg. No. 1258-723

ACCEPTED

Feb 18, 1967

INTENTION ACE

1258-723

