

566-551

6961-6 hww



Olin

CONTENTS 4 LBS. NET

SEVIN 5% DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 5.0%

INERT INGREDIENTS: 95.0%

TOTAL 100.0%

* Trademark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF INHALED SWALLOWED OR ABSORBED THROUGH SKIN. DO NOT BREATHE DUST DO NOT TAKE INTERNALLY AVOID PROLONGED OR REPEATED BREATHING OF DUST SKIN CONTACT MAY BE HARMFUL AVOID CONTACT WITH SKIN WASH HANDS BEFORE EATING TAKE SHOWER OR BATH AFTER HANDLING WEAR REGULAR LONG SLEEVED WORK CLOTHING CHANGE TO CLEAN CLOTHING DAILY AVOID CONTAMINATION OF FEED AND FOODSTUFFS

Note for Physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

USE PRECAUTIONS: Destroy empty bags by burning (KEEP OUT OF SMOKE).

For protection of honey bees avoid use, if possible, during periods when bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

GENERAL INFORMATION: Use a good type duster to obtain full coverage of foliage. Where a dosage range is indicated, use lower dosage rate on young plants; higher rate on mature plants. Apply when insects or their damage appear. Repeat at 7 to 10-day intervals or as necessary unless shorter range interval is specified.

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

SEVIN 5% DUST

DIRECTIONS FOR USE

BEANS (Green beans, lima beans, snap beans, cowpeas and blackeyed peas: Use 10 lbs. per acre for Mexican bean beetle; 20 lbs. per acre for bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle, leafhoppers, velvetbean caterpillar and Western bean cutworm. Use 20 to 30 lbs. for armyworms, corn earworm, stink bug and tarnished plant bug.

CUCUMBERS, MELONS, PUMPKIN and SQUASH: Use 10 to 20 lbs. per acre for pickleworm and melonworm; 20 lbs. for cucumber beetle, flea beetle, leafhopper and squash bug. **AVOID EXCESSIVE APPLICATIONS.** Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. **DO NOT USE ON WATERMELONS IN FLORIDA.**

EGGPLANT, PEPPER, POTATO and TOMATO: Use 10 to 20 lbs. per acre for Colorado potato beetle, flea beetle and leafhoppers. Use 20 to 40 lbs. per acre for European corn borer, fall armyworm, lace bug, tomato fruitworm, tomato hornworm, tarnished plant bug and stink bug.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER and KOHLRABI: Use 10 to 20 lbs. per acre for flea beetles and harlequin bug. Use 20 to 40 lbs. per acre for armyworms, cabbage caterpillars and corn earworm. Where cabbage looper is the principal problem, use an alternate pesticide that is specifically recommended. Do not apply within 3 days of harvest.

POULTRY (Chickens, Turkeys, Ducks, Geese, Game Birds, Pigeons): For control of Northern fowl mite, chicken mite, lice, fleas.

ON BIRDS: Dust birds thoroughly, using at the rate of 1 lb. 5% dust per 100 birds. **ON LITTER:** Apply at the rate of 1 lb. 5% dust per 40 square feet. Do not apply on or house birds in treated areas within 7 days of slaughter. Do not repeat applications within 4 weeks. Do not contaminate feed or drinking water.

DOGS and Cats: For control of brown dog tick and fleas. Rub thoroughly into hair or fur. Dust sleeping quarters weekly. Do not contaminate feed or drinking water. Do not apply to kittens under 4 weeks old.

TURF: For control of chinch bugs, fall armyworm, leafhoppers, sod webworm, ants, earwigs, fleas, millipedes and mosquitoes, dust lawn thoroughly at the rate of approximately 1 lb. 5% dust per 150 sq. ft. For best results, mow lawn, apply dust and water immediately. Repeat in 2 to 3 weeks if necessary.

LR356E-01R 6023952
USDA Reg. No. 1258-725

AGRICULTURAL DIVISION **Olin**
 LITTLE ROCK, ARKANSAS
 JOHN M. HILLISSION, CHEMICAL CORPORATION
 DALLAS, TEX. OMAHA, NEB. LITTLE ROCK, ARK.
 HOUSTON, TEX. PHOENIX, ARIZ. JACKSONVILLE, FLA.
 MOBILE, ALA. DULUTH, GA. WICHITA, KAN.
 SAVANNAH, MISS.

AGRICULTURAL DIVISION **Olin**
 LITTLE ROCK, ARKANSAS
 JOHN M. HILLISSION, CHEMICAL CORPORATION

lin
CONTENTS - LBS NET

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) **5.0%**

INERT INGREDIENTS: 95.0%

TOTAL 100.0%

Contains 5.0% Carbaryl (1-naphthyl N-methylcarbamate) and 95.0% inert ingredients.

USE PRECAUTIONS: Destroy empty bags by burning (KEEP OUT OF SMOKE). For protection of honey bees avoid use if possible during periods when bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

GENERAL INFORMATION: This insecticide is effective against a wide range of insects and mites. It is particularly effective against beetles, caterpillars, and other insects that feed on plants. It is also effective against some mites and ticks.

NOTICE TO BUYER

This product is registered under the Federal Insecticide, Fungicide, and Rodenticide Act. It is sold under the name of this manufacturer. The name of this manufacturer is printed on the label. The name of the manufacturer is also printed on the label. The name of the manufacturer is also printed on the label. The name of the manufacturer is also printed on the label.

BEANS: (Green beans, lima beans, snap beans, winged peas and brackeye peas) Use 10 lbs. per acre for Mexican bean beetle; 20 lbs. per acre for bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle, leafhoppers, velvetbean caterpillar and Western bean cutworm. Use 20 to 30 lbs. for armyworms, corn earworm, stink bug and tarnished plant bug.

CUCUMBERS, MELONS, PUMPKIN and SQUASH: Use 10 to 20 lbs. per acre for pickleworm and melonworm; 20 lbs. for cucumber beetle, flea beetle, leafhopper and squash bug. **AVOID EXCESSIVE APPLICATIONS.** Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. **DO NOT USE ON WATERMELONS IN FLORIDA.**

EGGPLANT, PEPPER, POTATO and TOMATO: Use 10 to 20 lbs. per acre for Colorado potato beetle, flea beetle and leafhoppers. Use 20 to 40 lbs. per acre for European corn borer, fall armyworm, lace bug, tomato fruitworm, tomato hornworm, tarnished plant bug and stink bug.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER and KOHLRABI: Use 10 to 20 lbs. per acre for flea beetles and harlequin bug. Use 20 to 40 lbs. per acre for armyworms, cabbage caterpillars and corn earworm. Where cabbage looper is the principal problem, use an alternate pesticide which is positively recommended. Do not apply within 3 days of harvest.

POULTRY: Chickens, Turkeys, Ducks, Geese, Game Birds, Pigeons. For control of Northern fowl mite, tickler mite, lice, etc. **ON BIRDS:** Dust birds thoroughly, using at the rate of 1/2 lb. of dust per 100 birds. **ON LITTER:** Apply at the rate of 1/2 lb. of dust per 40 square feet. Do not apply on or to birds in treated areas within 7 days of slaughter. Do not retreat applications within 4 weeks. Do not contaminate feed or drinking water.

DOGS and Cats: For control of ticks, fleas, and other parasites, use this product on the animal's body. Do not use on the face, ears, or eyes. Do not use on cats.

TURF: Use this product to control fleas, ticks, and other parasites on lawns. Apply at the rate of 1/2 lb. per 1000 square feet. Do not use on lawns that are to be seeded or sodded.