

1258-740

11/17/1965

1/2

OLIN MATHIESON CHEMICAL CORPORATION

745 FIFTH AVENUE

NEW YORK, NEW YORK 10022

ATTENTION: RICHARD F. PHILPITT

MANAGER, LEGISLATIVE & REGULATORY AFFAIRS

ACCEPTED

NOV 17 1965

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 1258-740


September 21, 1965

TERRACLOR SUPER X EMULSIFIABLE SOIL FUNGICIDE
TERRACLOR PLUS TERRAZOLE

FEDERAL REGISTRATION NUMBER 1258-740

NET CONTENTS: 55 GALLONS

OLIN
TERRACLOR(R) SUPER X EMULSIFIABLE
Soil Fungicide
Terraclor(R) Plus Terrazole*



Controls Rhizoctonia, Pythium, and Fusarium--Soil-borne seedling diseases of cotton.

ACTIVE INGREDIENTS:

| | |
|---|-------|
| Pentachloronitrobenzene..... | 23.0% |
| 5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole..... | 6.1% |
| INERT INGREDIENTS:..... | 70.9% |

| | |
|-------|--------|
| TOTAL | 100.0% |
|-------|--------|

Contains 2 lbs. of Pentachloronitrobenzene and 0.5 lb. of 5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole per U. S. gallon at 68° F.

USDA Reg. No. 1258-740

Terrazole*, Terraclor(R), Terraclor(R) Super X, and Olin are OLIN trademarks.

* Patent applied for.

WARNING (18 pt. caps): Do not inhale spray mist. Do not get on skin or in the eyes. May cause skin irritation. Wash thoroughly all exposed skin. In case of eye contact, flush with plenty of water. Repeated contact may be harmful. Wash thoroughly all contaminated clothing before reuse. Do not take internally. Do not store near heat or open flame. Use adequate ventilation. Keep container closed. Do not contaminate feed or food stuffs. Keep Out of the Reach of Children. (12 pt. type).

DIRECTIONS FOR USE ON COTTON (Seedling Disease Complex--Rhizoctonia, Pythium, and Fusarium): For soil in-furrow spray applications at planting time, use 2 - 3 qts. per acre mixed with 12 - 15 gallons of water. Spray all the soil that surrounds and covers the seed. Two nozzles work best; center the front nozzle over the open furrow to spray soil around the seed and the rear nozzle to spray covering soil as it fills the furrow.

COTTON SEED TREATMENT: Mix 8 - 12 oz. with 1 - 1½ pints of water and spray onto 100 lbs. of seed. Soil applications are required for later stages of pre-emergent and post-emergent damping-off. Treated seed should be labeled "Treated seed--do not use for food, feed or oil purposes."

NOTICE TO BUYER: No warranty of any kind, express or implied, is made concerning the use of this product. User assumes all risk and liability resulting from handling, use or application.

OLIN Agricultural Division
Commercial Development Department
745 Fifth Avenue, New York, N.Y. 10022
Olin Mathieson Chemical Corporation