

**Olin**

# Terraclor Super-X

**Soil Fungicide — EMULSIFIABLE**

1258-740  
**ACCEPTED**  
 10/2/72  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND ROVIDENTICIDE ACT  
 FOR ECONOMIC REGISTRATION  
 ED UNDER NO. 1258-740  
 ATTACHED TO THE

Controls Rhizoctonia, Pythium and Fusarium  
 — Soil-borne seedling diseases of cotton.

<b>ACTIVE INGREDIENTS:</b>	
Pentachloronitrobenzene	23.2%
5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole*	5.8%
<b>INERT INGREDIENTS:</b>	71.0%
*U.S. Patents 3,260,588 and 3,260,725; others pending. <b>TOTAL 100.0%</b>	
Contains 2 lbs. of Pentachloronitrobenzene and 0.5 lb. of 5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole per U. S. gallon at 68° F.	

## WARNING

**KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Do not inhale spray mist. Do not get on skin or in eyes. May cause skin irritation. Skin contact may be harmful. Wash thoroughly all exposed skin. In case of eye contact, flush with plenty of water. Wash thoroughly all contaminated clothing before reuse. Keep container closed. Do not contaminate feed of foodstuffs.

## CAUTION COMBUSTIBLE LIQUID.

**DO NOT USE, POUR, SPILL OR STORE  
 NEAR HEAT OR OPEN FLAME**

**DIRECTIONS FOR USE ON COTTON** (Seedling Disease Complex — Rhizoctonia, Pythium and Fusarium):

### SOIL IN-FURROW APPLICATION:

At planting time, use 2 to 3 qts. per acre (13,000 linear ft. based on 40 in. rows) mixed with 5 to 15 gallons of water. Spray all the soil that surrounds and covers the seed. Two nozzles work best; center the front nozzle over the open furrow to spray soil around the seed and the rear nozzle to spray the covering soil as it fills the furrow. Do not allow the feeding or grazing of cotton foliage by livestock. Do not feed gin waste to livestock.

### USE PRECAUTIONS

Do not plant treated fields to any crop other than cotton.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift of spray from the area being treated. Do not contaminate water by cleaning of equipment by disposal of wastes. Apply this product only as specified on this label.

**DESTROY THIS CONTAINER WHEN EMPTY — NEVER RE-USE**

Terrazole®, Terraclor®, and **Olin** are trademarks of Olin Corporation

### NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

LR347D-10V

**EPA Reg. No. 1258-740-AA**

AGRICULTURAL DIVISION  
 U.S. DEPARTMENT OF AGRICULTURE

**Olin**



# Terraclor Super-X

Soil Fungicide — EMULSIFIABLE

Controls Rhizoctonia, Pythium, and Fusarium

and some seedling diseases of cotton

## ACTIVE INGREDIENTS:

Pentachloronitrobenzene 23.2%

5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole 5.8%

INERT INGREDIENTS: 71.0%

Contains 2.4% of active ingredient TOTAL 100.0%

Contains 1.5% of Pentachloronitrobenzene and 0.5% of 5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole per gallon of emulsion

## WARNING

KEEP OUT OF REACH OF CHILDREN

Avoided. Do not get on skin or in eyes. If on skin, wash thoroughly with exposed skin. If in eyes, flush with plenty of water. Wash thoroughly and change clothing before use. Keep container closed. Do not contaminate feed or foodstuffs.

CAUTION COMBUSTIBLE LIQUID.

DO NOT USE, POUR, SPILL OR STORE  
NEAR HEAT OR OPEN FLAME

DIRECTIONS FOR USE ON COTTON (Seedling Disease Complex Rhizoctonia, Pythium and Fusarium)

### SOIL IN-FURROW APPLICATION:

At planting time, use 2 to 3 gts. per acre (13,000 linear ft. banded on 40 in. rows) mixed with 5 to 15 gallons of water. Spray all the soil that furrows and covers the seed. Two nozzles work best. Center the front nozzle over the inter-furrow to spray soil around the seed and the rear nozzle to spray the covering soil as it fills the furrow. Do not allow the feeding or grazing of cotton foliage by livestock or wildlife.

### USE PRECAUTIONS

Do not plant treated fields to any crop other than cotton.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift of spray from the area being treated. Do not contaminate water by cleaning of equipment by disposal of wastes. Apply this product only as specified on this label.

DESTROY THIS CONTAINER WHEN EMPTY NEVER RE-USE

Olin

### NOTICE TO BUYER

This product is a registered pesticide of the U.S. Department of Agriculture. It may be used only on the crops, plants and animals listed on its label. It must be used in the manner and at the rate of application specified on its label. It is unlawful to use this product in any manner inconsistent with the label. The U.S. Government is authorized to reproduce and distribute reprints for government purposes not withstanding any copyright notation that may appear hereon.

4347E-10V

EPA Reg. No. 1258-740-AA

AGRICULTURAL DIVISION  
LITTLE ROCK, ARKANSAS  
OLIN CORPORATION

