

TURFCIDE 10 GRANULAR

0% ACCEPTED
1-15-68

1258-765

SOIL FUNGICIDE
CONTROLS BROWN PATCH OF CERTAIN TURF

ACTIVE INGREDIENTS:

Pentachloronitrobenzene 10%

INERT INGREDIENTS: 90%

Total 100%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid inhaling dust. This material may cause skin irritation. Wash hands and face thoroughly after using. Wear gloves and a respirator when loading the spreader or duster. Do not contaminate feed or foodstuffs.

NOTICE TO BUYER

Olin warrants that this product conforms to the description on this label and is reasonably fit for use as directed hereon. Olin neither makes nor authorizes any representative, agent or other person to make any other warranty or representation, express or implied, with respect to this product. Because unforeseeable factors beyond Olin's control prevent it from eliminating all risks of use of chemical products, even though reasonably fit for such use, this product is sold upon condition that the buyer and user assume all risks and liability, except as stated in sentence one, resulting from the storage, handling or use of this product, including, without limitation, damage to plants, crops and animals to which the product is applied, failure to control pests, compatibility with other products, drift, contamination of equipment or personal injury.

DIRECTIONS FOR USE

LAWNS (Brown Patch — Rhizoctonia solani): For Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses: Make one application in the fall or spring at first indication of infection. Apply 7½ lbs. per 1,000 sq. ft. Following application, the treated area should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3 to 4 weeks later. For Dichondra: Treat as above using 7½ lbs. per 1,000 square feet.

For Northern or cool season lawns such as Rye Grass, Fescue, Bluegrass, Bent and similar grasses: Apply 2 to 2½ lbs. per 1,000 sq. ft. at first sign of disease. Repeat every 7 to 10 days during warm, damp weather. Water treated area lightly to wash chemical down to soil.

Note: Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

LR406A-03Q 6024127
USDA Reg. No. 1258-765

TURFCIDE and OLIN are OMCC trademarks

SAVAGE, MINN.
BALTIMORE, MD.
HOUSTON, TEX.
NO. LITTLE ROCK, ARK.
OMAHA, NEB.

AGRICULTURAL DIVISION
LITTLE ROCK, ARKANSAS



PHOENIX, ARIZ.
RICHMOND, IND.
ST. LOUIS, MO.
WILLIAMSTON, N.C.
WICHITA, KAN.

OLIN MATHIESON CHEMICAL CORPORATION

1258-765

01/15/1968

1/1