

3"

10"


1-5/8"

7/16" TREATS OVER 2,650 SQUARE FEET

10% GRANULAR  
FUNGICIDE 7/16"

2-3/16" 3/4" Turfcide®  
10% GRANULAR  
FUNGICIDE 3/4"

26"



NOTE: THESE DIMENSIONS ARE BASED ON A 10" X 3" X 26"  
 OPEN-MOUTH PAPER BAG SIZE. THIS MINIATURE ART IS IN  
 APPROXIMATE PROPORTION AND SHOULD ENLARGE ACCORDINGLY.  
 PLEASE ADHERE TO TYPE STYLES AND POSITIONS USED ON MINIATURE.

10"



1-5/8"



**10% GRANULAR FUNGICIDE**

7/16"

7/16"

**DIRECTIONS**

Lawns—To control Brown Patch (*Rhizoctonia Solani*)—For Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses make one application in the fall or spring at first indication of infection.

Apply 7½ pounds per 1,000 square feet. Following application the treated area should be lightly watered to insure movement through the grass to the soil level. If the treated area is subjected to unusually heavy rainfall or is flooded, or if the disease is severe and reappears, the area should be retreated 3 to 4 weeks later. For Dichondra: Treat as above using 7½ pounds per 1,000 square feet.

Note — Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**  
AVOID INHALING DUST. THIS MATERIAL MAY CAUSE SKIN IRRITATION. WASH HANDS AND FACE THOROUGHLY AFTER USING. WEAR GLOVES AND A RESPIRATOR WHEN LOADING THE SPREADER OR DUSTER.

DO NOT CONTAMINATE FEED OR FOODSTUFFS. THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF ANY BODY OF WATER. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.

**NOTICE TO BUYER**

Seller's attention shall be limited to the terms of the

26"

## DIRECTIONS

Lawns—To control Brown Patch (*Rhizoctonia Solani*)—  
For Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses make one application in the fall or spring at first indication of infection.

Apply  $7\frac{1}{2}$  pounds per 1,000 square feet. Following application the treated area should be lightly watered to insure movement through the grass to the soil level. If the treated area is subjected to unusually heavy rainfall or is flooded, or if the disease is severe and reappears, the area should be retreated 3 to 4 weeks later. For Dichondra: Treat as above using  $7\frac{1}{2}$  pounds per 1,000 square feet.

Note — Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

AVOID INHALING DUST. THIS MATERIAL MAY CAUSE SKIN IRRITATION. WASH HANDS AND FACE THOROUGHLY AFTER USING. WEAR GLOVES AND A RESPIRATOR WHEN LOADING THE SPREADER OR DUSTER.

DO NOT CONTAMINATE FEED OR FOODSTUFFS. THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF ANY BODY OF WATER. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.

## NOTICE TO BUYER

Seller's guarantee shall be limited to the terms of the label and subject thereto the Buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

TURFIDE and Olin are trademarks

NET WEIGHT 20 POUNDS

AGRICULTURAL DIVISION

LITTLE ROCK, ARKANSAS  
OLIN CORPORATION

**Olin**

**EPA**

USDA Reg. No. 1258-765

LR 406A-03Q