

NET WEIGHT
20 POUNDS

Turficide®

10% GRANULAR
SOIL FUNGICIDE

Controls Certain Soil-Borne
Diseases of Lawns — Turf — Ornamentals

ACTIVE INGREDIENT:

Penclorolol nitrobenzoyl

INERT INGREDIENTS:

Total

10%
90%
100%

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT
(First Aid Procedures)

If swallowed, induce vomiting. If on skin, remove contaminated clothing and wash skin with soap and warm water. If inhaled, remove to fresh air. If in eyes, flush with water for 15 minutes.

For further treatment, get medical attention. See rear panel for additional precautionary statements.

Turficide® is a trademark of Olin Corporation.

EPA Reg No 1258-765

EPA Est 1258 MS-1



P.O. Box 991
Little Rock, AR 72203

10% GRANULAR
SOIL FUNGICIDE

Turficide®

ACCEPTED
JAN 8 1982
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the fungicide registered under EPA Reg. No. 1258-765

Turficide®
10% GRANULAR
SOIL FUNGICIDE

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. This material may cause skin irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt while handling product. Wash hands and face after use and prior to eating, drinking or smoking. Wash contaminated clothing before reuse. Do not handle or apply product with bare hands. Keep out of reach of children.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply product in areas which would permit runoff to contaminate water supplies.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

LAWNS AND TURF

Brown patch—Rhizoctonia solani: For Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses. Make one application in the fall or spring at the rate of 1.0 lb. per 1,000 sq. ft. Following applications, water the lawn thoroughly. If the lawn is heavily infested, a second application may be made. If the treated area is subjected to unusually heavy rainfall or flooded, or if the disease is severe or reappears, the area should be retreated 3 to 4 weeks later. For the best results, apply Turficide to the lawn in the fall or early spring. For cool season lawns such as Ryegrass, Fescue, Bluegrass, Bent or similar grasses, apply 2 to 2.5 lbs. per 1,000 sq. ft. at first sign of disease. Repeat every 7 to 10 days during warm, humid weather. When the weather turns cool, the grasses may naturally recover. If the disease is severe or reappears, the area should be retreated 3 to 4 weeks later.

Hemiphusium leaf spot: Apply at the rate of 5 to 7.5 lbs. per 1,000 sq. ft. Preventative applications should be made in spring or fall of each year. For control of existing infections, apply at the higher rate as soon as symptoms are first noted. Following applications, the treated area should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall or flooded, or if disease is severe or reappears, the area should be retreated 3 to 4 weeks later.

Snow mold (Typhula incarnata, Fusarium nivale): Apply at the rate of 5 to 7.5 lbs. per 1,000 sq. ft. in the fall or early spring. Water the lawn thoroughly after application. Application should be made prior to first snowfall.

Do not apply Turficide to areas of turf cuttings to livestock.

ORNAMENTALS

AFRICAN VIOLETS, BEGONIAS, CARNATIONS, CHRYSANTHEMUMS, POINSETTIAS, SNAPDRAGONS, and similar bedding plants (Stem Rot—Rhizoctonia): Apply 1.5 lbs. per 1,000 sq. ft. by spreading evenly on the soil surface and thoroughly mixing into the top 2 inches of soil. Apply to a well prepared seed bed.

BULBOUS IRIS, HYACINTH, NARCISSUS, and TULIP (Crown Rot and Black Rot): Apply broadcast 1.000 to 2.000 lbs. per acre (2 to 4 lbs. per 100 sq. ft.) and mix thoroughly into the top 2 inches of soil. Apply to the soil surface and mix with the soil. Do not use for heavier soils and for Black Rot control.

CALENDULA, LARKSPUR, SNAPDRAGON, and SWEET PEA (Stem Rot—Sclerotinia): Apply 1.000 lbs. per acre broadcast one week prior to planting. For smaller areas, apply 1 lb. per 100 sq. ft. Spread on the soil surface and thoroughly mix into the top 2 inches of soil. Turficide was developed for this use in Charleston, South Carolina.

DUTCH IRIS (Bulb and stem rot caused by Sclerotinia rot): Broadcast Apply 1.000 to 1.500 lbs. per acre (2 to 4 lbs. per 100 sq. ft.) prior to planting and mix with the top 2 inches of soil by discing or rototilling.

EASTERN LILIES (Rhizoctonia root rot): Broadcast Apply 1.000 to 2.000 lbs. per acre (2 to 4 lbs. per 100 sq. ft.) to soil surface prior to planting and mix thoroughly to upper 2 inches by double discing or rototilling. Note: Use the lowest rate of application for heavy soils and higher rate for applications in heavier soils.

FLORIST'S HYDRANGEA (Stem and leaf blight caused by Botrytis blight): Apply 1.000 to 1.500 lbs. per acre (2 to 4 lbs. per 100 sq. ft.) to soil surface and mix with the soil. Apply thoroughly in upper 4 to 6 inches by double discing or rototilling. (Bulbs may be planted immediately.)

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Dispose of bags according to approved Federal, State or local procedures under the Resource and Recovery Act.

GENERAL: Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.