

# 

20-5 SEED DUST FUNGICIDE

Controls Soil-borne seedling diseases of cotton.

#### **ACTIVE INGREDIENTS:**

Pentachloronitrobenzene 20.0% 5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole\* 5.0% INERT INGREDIENTS: 75.0% TOTAL 100.0%

# WARNING

# KEEP OUT OF REACH OF CHILDREN

The second of th

# PRECAUTIONS IN USING

Rinse equipment and dispose of wastes by burying in non-crop land away from water supplies. Empty bags should be disposed of by burying with wastes or burning (KEEP OUT OF SMOKE).

Treated seud must not be used for feed, food, or oil purposes.

## **DIRECTIONS FOR USE**

### COTTON:

PLANTER BOX TREATMENT: (Seedling Disease Complex-Rhizoctonia, Pythium, Thielaviopsis and Fusarium).

Mix 1 to 1½ lbs. of Terraclor Super-X 20-5 Dust with the quantity of cottonseed to be planted to one acre (per 13,000 linear feet based on 40 inch rows) and apply at planting time through planter box.

## NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

Terraclor Super-X,® Terraclor,® Terrazole,® and Olin are trademarks of OMCC.

\*U S Patents 3,260,588 and 3,260,725; others pending

LR455-12R USDA Reg. No. 1258-813