

Important — Federal law requires that treated seed be dyed to impart an unnatural color and that bags containing treated seed shall be labeled with the following information:

This seed has been treated
with
Pentachloronitrobenzene
and
5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole

CAUTION
DO NOT USE FOR FEED, FOOD OR OIL PURPOSES.
STORE AWAY FROM FEEDS AND FOODSTUFFS

USE CAUTIONS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

DESTROY THIS CONTAINER WHEN EMPTY
— NEVER REUSE.

COMBUSTIBLE LIQUID

DO NOT USE, POUR, SPILL OR STORE
NEAR HEAT OR OPEN FLAME.

Keep container closed. Use with adequate ventilation.

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

Terra-Coat,® Terraclor,® Terrazole,® and Olin are
trademarks of Olin Corporation

Terra-Coat[®]

L-205

SEED TREATMENT FUNGICIDE

Terraclor[®] Plus Terrazole[®]
WITH DYE

ACTIVE INGREDIENTS:

Pentachloronitrobenzene 23.2%

5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole* 5.8%

INERT INGREDIENTS: 71.0%

TOTAL 100.0%

Contains 2 lbs. of Pentachloronitrobenzene and 0.5 lb. of 5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole per U. S. gallon at 68° F.

*U. S. Patents 3,260,588 and 3,260,725; others pending.

WARNING

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. Do not inhale spray mist. Do not get on skin, on clothing or in eyes. May cause skin irritation. Wash thoroughly all exposed skin. Contact may be harmful. Wash thoroughly all contaminated clothing before reuse. In case of eye contact, flush with plenty of water. Do not contaminate feed or foodstuffs.

EPA Reg. No. 1258-818-AA

PROTECT FROM TEMPERATURES BELOW 20° F.

AGRICULTURAL DIVISION **Olin**
LITTLE ROCK, ARKANSAS
ON

DIRECTIONS FOR USE

Controls certain seed and soil-borne seedling diseases. Apply Terra-Coat L-205 undiluted directly to seed (or mix with one to two parts of water as required for optimum seed coverage) at dosage rates indicated in the chart below. This product contains a dye to indicate treated seed.

Planting Seeds	Application Rates in fluid ounces	Seedling Disease Complex
Beans	2 oz./bushel	Rhizoctonia, Phythium, Thielaviopsis and Fusarium
Corn	2 oz./100 lbs.	
Cotton	12-15 oz./100 lbs.	
*Peanuts	2-4 oz./100 lbs.	
Peas	2-4 oz./bushel	
Sorghum	2 oz./100 lbs.	
Soybeans	2-4 oz./bushel	Seed-borne rust, Rhizoctonia, Phythium, Thielaviopsis and Fusarium
Rice	2-4 oz./bushel	
Safflower	2-4 oz./100 lbs.	
Sugar Beet	6-12 oz./100 lbs.	Aphanomyces, Rhizoctonia, Phythium, Thielaviopsis and Fusarium
Wheat	2 oz./bushel	Common Smut or Bunt, Rhizoctonia, Phythium, Thielaviopsis and Fusarium
Barley	2-4 oz./bushel	Covered Smut

*Note: In the treatment of Peanuts, do not dilute with water.

LR460C-12U

6324950

