

ACTIVE INGREDIENT:

Pentachloronitrobenzene

INERT INGREDIENTS:

Total - -----

Terraclor[®] is a trademark of Olin Corporation.

CAUTION **KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Avoid inhaling dust. This material may cause skin irritation. Wash hands and face thoroughly after using. Wear gloves and a respirator when loading the spreader Do not contaminate feed or foodstuffs.

USE CAUTIONS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil.

Destroy empty bags by burning (KEEP OUT OF SMOKE).

DIRECTIONS FOR USE

CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, AND BROCCOLI (Clubroot - Plasmodiophora brassicae): Row Application Apply 67 by per 13,100 feet of row for cabbage. preccol, and brussely shouts or to 10,900 mean feet of row for Lauthower in a 12 to 15 it to basid, by means of a suitable applicator, and rotote to a depth of 4 to 6 increasionm eduate i, prior to transr lanting

CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, AND BROCCOLI (Clubroot-Plasmodiophora brassicae):

BROADCAST Apply the use user acre to the sole surface prove In transplanting Disclard and into its proladepth of 4 to elimenes The TELE in disposed if shares a period were three on and the Same prouts at sivery important to the ipport of peritor to the

CAULIFLOWER (Wire Stem or Blank root - Corticium solani): Abd 2 3 1 more de la 2 en en anter and the second second

70% 100% NOV 28 1972

UNDER THE PEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER-ED UNDER NO. 2012 SUBJECT TO ATTACHED COMMENTS.

BEANS (Root and Stem Rot - Rhizoctonia Solani): FURROW TREATMENT:- Apply 4 to 5 lbs. per 14,500 linear feet of row for bush beans, or per 8,430 linear feet of row for pole beans. Direct the material into the planting furrow and covering soil at planting time. Avoid getting the material concentrated around the seed otherwise delayed emergence may occur. Do not feed treated vines to livestock.

pegging zone.

a depth of 4 to 6 inches. ROW (in-furrow) APPLICATION: Apply 33 lbs. per 12,400 linear feet of row and work thoroughly into the top 4 to 6 inches of soil (it is essential that the fungicide is worked into the tuberforming zone of the soil). It is desirable to make treatment when the soil is slightly moist.

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

30% ACCEPTED Controls Certain Soll-Borne Diseases Of BEANS CRUCIFERS POTATOES PEANUTS •

EPA Reg. No. 1258-823-AA LR458D-02V

DIRECTIONS FOR USE (Con't.)

PEANUTS (Southern Stem and Root Rot - Sclerotium Rolfsii): PRE-PLANT SOIL MIX:- Apply 34 lbs. per 12,400 linear feet of row, in an 8 to 12 inch band, centered on the row and mix to a depth of 2 to 4 inches. Application may be made simultaneously with planting, using suitable equipment.

SURFACE BAND TREATMENT: - When pegging begins, apply 34 lbs. per 12,400 linear feet of row to an 8 to 12 inch band, directed at the plant crown and soil

Note: Do not use more than one application method during the same planting season. Do not feed treated vines to livestock.

POTATOES (Rhizoctonia solani): BROADCAST TREATMENT:- Use 60 to 83 lbs. per acre. Spread evenly over the soil surface prior to planting. Disc and cross disc to

NOTICE TO BUYER