

Clin

Terrazole

35% WETTABLE POWDER

controls certain soil-borne diseases of
ORNAMENTAL TURF and ORNAMENTALS

ACTIVE INGREDIENTS:

5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole*

35%

INERT INGREDIENTS:

65%

TOTAL

100%

*U. S. Patents 3,260,588 and 3,260,725; others pending.

WARNING

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION.

Avoid inhalation of dust and spray mist.

Wash hands and face thoroughly after using.

Avoid contamination of feed and foodstuffs.

USE CAUTIONS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Empty bags should be disposed of by burying with wastes or burning (KEEP OUT OF SMOKE).

EPA Reg. No. 1258-826-AA

25
LBS. NET

AGRICULTURAL DIVISION

Clin

LITTLE ROCK, ARKANSAS

CLIN CORPORATION

GENERAL INFORMATION

Terrazole 35% WP is useful as a soil fungicide for control of damping-off, root and stem diseases caused by Pythium and Phytophthora.

Use only at recommended rates as overdosage may be harmful to sensitive plants. Do not mix with other chemicals, other than Terraclor^(R), unless prior use has proven compatibility.

DO NOT PLANT FOOD CROPS IN ROTATION ON TREATED SOILS.

BEDDING PLANTS

<u>Ageratum</u>	<u>Coleus</u>	<u>Petunia</u>
<u>Aster</u>	<u>Dahlia</u>	<u>Portulaca</u>
<u>Balsam</u>	<u>Dianthus</u>	<u>Salvia</u>
<u>Begonia</u>	<u>Foxglove</u>	<u>Stokesia</u>
<u>Calendula</u>	<u>Impatiens</u>	<u>Verbena</u>
<u>Carnation</u>	<u>Marigold</u>	<u>Vinca</u>
<u>Celosia</u>	<u>Pansy</u>	<u>Zinnia</u>
	<u>Snapdragon</u>	

TROPICAL FOLIAGE PLANTS

<u>Aglaonema</u>	<u>Nephtytis</u>
<u>Anthurium</u>	<u>Peperomia</u>
<u>Chinese Evergreen</u>	<u>Pothos</u>
<u>Dieffenbachia</u>	<u>Pilea</u>
	<u>Philodendron Spp.</u>

HERBACIOUS and WOODY PLANTS

<u>Chrysanthemum</u>	<u>Geonothus</u>
<u>Easter Lily</u>	<u>Poinsettia</u>
<u>Arctostaphylus</u>	<u>Rhamnus</u>

RECOMMENDATIONS

For bedding and bench plantings use Terrazole 35% WP mixed with water, as a soil drench at time of seeding and transplanting. The drench solution should be continually agitated to assure uniform application of the fungicidal material. To improve penetration of the fungicide, the drenching should be followed with an equal quantity of water, applied either by sprinkling or by irrigation.

BEDDING PLANTS (soil media 2" to 3" deep)—Add 2 to 3 ozs. to 50 gallons of water and apply to 400 sq. ft. of bed area. Repeat applications may be made at one to two month intervals, if necessary.

TROPICAL FOLIAGE PLANTS (grown in peat or other high organic media, 6" deep).

BENCH PLANTINGS—Add 4 to 6 ozs. to 50 gallons of water and apply to 200 sq. ft. of bench area.

BEDGROWN and POT PLANTINGS—Add 1½ to 5 ozs. in 50 gallons of water. For bed areas apply to 200 sq. ft. of soil mixture or in sufficient amount to saturate the soil.

For pots use approximately ½ pint solution per 6" container.

Repeat applications at 2 to 3 month intervals if required.

DRY SOIL MIX—For plants listed above, blend 1½ to 3 ozs. Terrazole into one cubic yard of soil mixture. Blending may be improved by first premixing the fungicide with five or six parts of soil and then blending into the soil mixture.

ORNAMENTAL TURF—Pythium fungi (cottony blight, grease spot, damping-off). Mix 4 to 8 ozs. in 5 gals. of water and apply to 1,000 sq. ft. of established ornamental turf. Begin applications at first sign of disease and retreat in 5 to 10 days depending upon climatic conditions to maintain disease control.

On newly seeded areas (damping-off caused by species of Pythium fungi) mix 6 to 8 ozs. in 5 to 10 gallons of water and apply to 1,000 sq. ft. of area immediately after seeding. retreat at 5 to 7 day intervals to maintain a preventive disease control program.

Do not allow animals to graze on treated areas. Do not feed clippings from treated areas to animals.

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions or under abnormal conditions, or in combination with other materials or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

Terrazole[®] and Clin are trademarks of Clin Corporation.

R474A 120