



\$1.65
Suggested
Retail Price

★ ★ ★ ★ CONTENTS 4 LBS. NET ★ ★ ★ ★

5-6

INSECTICIDE-FUNGICIDE GARDEN DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)..... 5%

Copper as metallic (from Tribasic Copper Sulphate)..... 4%

INERT INGREDIENTS:..... 91%

Total..... 100%

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF INHALED, SWALLOWED OR ABSORBED THROUGH SKIN. DO NOT BREATHE DUST. SKIN CONTACT MAY BE HARMFUL. AVOID CONTACT WITH SKIN. WASH HANDS BEFORE EATING. TAKE SHOWER OR BATH AFTER HANDLING. WEAR REGULAR LONG SLEEVED WORK CLOTHING. CHANGE TO CLEAN CLOTHING DAILY. AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

For protection of honey bees avoid use, if possible, during periods when bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

Destroy empty container by burning (KEEP OUT OF SMOKE).

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to the chemical description and is reasonably fit for the purpose stated on the label when used in accordance with the directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions or under abnormal conditions or in combination with other materials or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

AGRICULTURAL DIVISION **Olin**
GREEN ROCK, ARKANSAS
OLIN CORPORATION

★ ★ ★ ★ ★ ★ ★ ★ ★ ★ ★ ★ ★ ★ ★ ★

GENERAL INFORMATION

This is a combination insecticide-fungicide for use when certain insects and such diseases as Anthracnose, Downy Mildew, Septoria Leaf Spot, Nailhead rust, Early and Late Blight are a problem. These diseases are more easily prevented than eradicated after infection is apparent. Start applications when diseases have been reported in the area or when weather conditions are favorable for development. If insect control, however, is the only objective, revert to the use of SEVIN® 5% DUST. If insects are not present use a straight fungicide such as SPERLOX-Z Lawn & Garden Fungicide.

Repeat applications at 5, 7 or 10 day intervals depending upon the severity of disease and cool, moist weather conditions favorable to disease infection.

Dusters should be adjusted to give uniform and thorough coverage of foliage. Where a dosage range is indicated, use the lower rate on young plants, gradually increasing the rate to the maximum range for mature plants.

DIRECTIONS FOR USE

✓ **BEANS** (Green beans, lima beans, snap beans, cowpeas and black-eyed peas): Use 10 lbs. per acre for Mexican bean beetle; 20 lbs. per acre for bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle, leafhoppers, velvetbean caterpillar and Western bean cutworm. Use 20 to 30 lbs. for armyworms, corn earworm, stink bug and tarnished plant bug. Use 25 to 30 lbs. per acre on beans and lima beans to control anthracnose, downy and powdery mildew, and bean mildew.

✓ **CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER:** Use 10 to 20 lbs. per acre for flea beetles and harlequin bug; Use 20 to 40 lbs. per acre for armyworms, cabbage caterpillars and corn earworm. Where cabbage looper is the principal problem, use an alternate pesticide that is specifically recommended. Do not apply within 3 days of harvest. Use 25 to 35 lbs. per acre to control downy mildew and leaf spots.

CUCUMBERS, MELONS, PUMPKIN and SQUASH: Use 10 to 20 lbs. per acre for pickleworm and melonworm; 20 lbs. for cucumber beetle, flea beetle, leafhopper and squash bug. AVOID EXCESSIVE APPLICATIONS. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. DO NOT USE ON WATERMELONS IN FLORIDA. Use 25 to 40 lbs. to control anthracnose, downy and powdery mildew, and powdery mildew.

TOMATO, EGGPLANT, PEPPER, and POTATOES: Use 10 to 20 lbs. per acre for Colorado potato beetle, flea beetle and leafhoppers. Use 20 to 40 lbs. per acre for European corn borer, fall armyworm, lace bug, tomato fruitworm, tomato hornworm, tarnished plant bug and stink bug. Use 25 to 40 lbs. per acre to control nailhead rust on tomatoes and early and late blight on tomatoes and potatoes; 35 to 40 lbs. per acre on eggplants to control anthracnose, downy mildew and phomopsis blight; 35 to 40 lbs. per acre on peppers to control anthracnose, downy mildew and cercospora leafspot.

* Trademark of Union Carbide Corporation for the active ingredients, 1-naphthyl N-methylcarbamate.

6023635 12476C-020
USDA Reg. No. 1258-836

THIS CONTAINER IS PACKED BY WEIGHT AND
NOT BY VOLUME. IT WILL NOT APPEAR FULL

BEST DOCUMENT AVAILABLE

Accepted
5/22/77
1255-126



\$1.65
Suggested
Retail Price

★ ★ ★ ★ CONTENTS 4 LBS. NET ★ ★ ★ ★

5-6

**INSECTICIDE-FUNGICIDE
GARDEN DUST**

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)	5%
Copper as metallic (from Tribasic Copper Sulphate)	6%

INERT INGREDIENTS:

Total	89%
	100%

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed. Do not breathe dust. May be harmful if in contact with skin. Wash hands thoroughly after use. Wear regular long sleeves and pants. Change clothes after use. Avoid contamination of food and feed stuffs.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

For protection of honey bees avoid use, if possible, during periods when bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

Destroy empty container by burning (KEEP OUT OF SMOKE)

NOTICE TO BUYER

This product is a registered trademark of Olin Corporation. It is sold as a general purpose insecticide and fungicide. It is not to be used for any other purpose. It is not to be used in any other manner than as directed. It is not to be used in any other manner than as directed. It is not to be used in any other manner than as directed.

AGRICULTURAL DIVISION
LITTLE ROCK, ARKANSAS
OLIN CORPORATION

BEAN
10 lbs
ber be
Weste
and ta
beans

CABB
Use 10
acre fo
is the p
Do not
downy

CUCUM
picklev
squash
treatme
high hu
to co

TOMAT
Colora
Europe
tarnish
nailho
potato
downy
to co

The Jemere
N-methy co

INSECTICIDE-FUNGICIDE GARDEN DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 5%

Copper as metallic (from Tribasic Copper Sulphate) 6%

INERT INGREDIENTS: 89%

Total 100%

CAUTION KEEP OUT OF REACH OF CHILDREN

HARMFUL IF INHALED, SWALLOWED OR ABSORBED THROUGH SKIN. DO NOT BREATHE DUST. SKIN CONTACT MAY BE HARMFUL. AVOID CONTACT WITH SKIN. WASH HANDS BEFORE EATING. TAKE SHOWER OR BATH AFTER HANDLING. WEAR REGULAR LONG SLEEVED WORK CLOTHING. CHANGE TO CLEAN CLOTHING DAILY. AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

For protection of honey bees avoid use, if possible, during periods when bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

Destroy empty container by burning (KEEP OUT OF SMOKE).

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

BEANS (Green beans, lima beans, etc.)
Use 10 lbs. per acre for Mexican bean beetle, flea beetle, Japanese beetle, and tarnished plant bug. Use 10 to 20 lbs. per acre for Western bean cutworm. Use 10 to 20 lbs. per acre for beans to control anthracnose.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS
Use 10 to 20 lbs. per acre for armyworms, cabbage looper, and cabbage root fly. If downy mildew and leaf miner is the principal problem, use 10 to 20 lbs. per acre. Do not apply within 3 days of harvest.

CUCUMBERS, MELONS, PEPPERS
Use 10 to 20 lbs. per acre for pickleworm and melonworm. AVOID EXCESSIVE treatments are made when high humidity. DO NOT USE for control anthracnose.

TOMATO. EGGPLANT, PEPPERS
Use 10 to 20 lbs. per acre for Colorado potato beetle, flea beetle, European corn borer, fall armyworm, and tarnished plant bug and stem borer. Use 10 to 20 lbs. per acre for potatoes; 35 to 40 lbs. per acre for tomatoes.