13"

AMMO-PHOS BUILT 18/2-2 IN 12-12-12

With Dacthal[®] Pre-emergence Herbicide Added

MINIMUM GUARANTEED ANA

Total Nitrogen (N)	12%
Avail. Phos. Acid (P2O5)	
Soluble Potash (K ₂ O)	
Sulfur-Combined (S)	

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. AVOID CONTACT WITH EYES. WASH AFTER USING. AVOID CONTAMINATION OF FEED OR FOODSTUFFS. BURY OR BURN EMPTY BAGS AND STAY OUT OF SMOKE.

DIRECTIONS USE INFORMATION


Ammo-Phos Lawn Fertilizer with Dacthal Herbicide is effective on mineral soils in feeding lawn turfs and in the control of crabgrass, chickweed, carpetweed, green and yellow foxtail, goosegrass, lambsquarter, lovegrass, purslane, Texas millet and witchgrass. It should be applied before seeds germinate to be effective.

HOW TO APPLY: To control crabgrass and similar annual weeds, apply Ammo-Phos Lawn Fertilizer with Dacthal Herbicide at the rate of 20 pounds per 1,000 square feet in early spring prior to seed germination. Use a fertilizer spreader and apply uniformly. A second application at heli the above rate may be made two months later, if needed.

Remove leaves, tall or dead grass and other debris before applying. Do not disturb the soil surface after applying. Water thoroughly after applying. To control Poa Annua and other erratic late-germinating weeds, apply a supplementary treatment at the rate of 30 lbs. per 1,000 square feet in late summer or early full.

If reseeding is needed, delay at least 60 days from application.





per 1,000 square feet in early spring prior to seed germin fertilizer spreader and apply uniformly. A second application above rate may be made two months later, if needed.

Remove leaves, tall or dead gross and other debris before not disturb the soil surface after applying. Water thoroughly a To control Poa Annua and other erratic late-germinating we supplementary treatment at the rate of 30 lbs. per 1,000 squar summer or early fall.

If reseeding is needed, delay at least 60 days from applicati



SPREADER SETTINGS

Once Over Twice Over 6 5 25

Drop-2 pr. Spreadles

Numerous drop-type spreaders a Each has a different flow rate Don' proper setting. Follow these four si calibrate your spreader for AMMO-Fertilizer:

Measure two pounds (approxpints) of AMMO-PHOS # Lawn Fiplace it in spreader to check for 2 1,000 square feet application rate.
Determine distance spreader mucover 100 square feet by dividing spinto 1200. For example, if yours 60

wide sprea 'er, 20 1200 your tes 69 feet.

3. Measure a 60-foot length of clean sidewalk or drivewa spreader to an arbitrary setting and cover that distance. If the not available, cover 30 feet of sidewalk twice or 15 feet four 4. All material should be used in the 60 feet, regardless o rate per 1,000 square feet. If material remains after test run, ch number setting. If you use all material before test is complete smaller number setting and try again.

CAUTIONS

Keep children and pets off treated areas until material is soil and turf is completely dry. Do not use on dichondra a lawns. Do not graze livestock on treated area.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms of the label thereto the buyer assumes any risk to persons or property a the use or handling and accepts this product on these condition

Amma Phase is a trademark of Ofin Corporation Dorthath is a trademark of Diamond Shall each Corporation

NET WEIGHT 50 POUND.

AGRICULTURAL DIVISION

OLIN CORPORATION

LR480A-02U

USDA Reg.

nation. Use a n at half the	
applying. Do fter applying.	
eds, apply a re feet in late	32,
ion.	n
Spread Width 612 Feet	
ire available 't guess at the imple steps to	
-PHOSE Lawn ximately two ertilizer and	
20 pounds per ust operate to	
preader width is a 20-inch- t run will be	
y. Open the at distance is times. of application	
noose a larger red, choose a	
watered into and bentgrass	
l and subject trising out of ons.	
n	
	•
No. 1258-864	