



Terrazole 25% EMULSIFIABLE

controls certain soil-borne diseases
of ORNAMENTALS

ACTIVE INGREDIENT	
5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole*	25.0%
INERT INGREDIENTS:	75.0%
TOTAL	100.0%

*U. S. Patents 3,260,588 and 3,260,725; others pending.

WARNING

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not inhale spray mist. Do not get on skin or in eyes. May cause skin irritation. Skin contact may be harmful. Wash thoroughly all exposed skin. In case of eye contact, flush with plenty of water. Wash thoroughly all contaminated clothing before reuse.

EPA Reg. No. 1258-931-AA

ONE QUART

CAUTION: COMBUSTIBLE LIQUID

Keep away from heat and open flame.

LR491A-06V

AGRICULTURAL DIVISION

11111 ROCK AVE

OLIN CORPORATION



931

7/17/70
UNDER THE FEDERAL IN-
FUNGICIDE ACT AND REGU-
LATION FOR FERTILIZERS
ED UNDER 259.931

GENERAL INFORMATION

Terrazole is useful as a soil fungicide for control of damping-off, root and stem diseases caused by Pythium and Phytophthora. Use only at recommended rates as overdosage may be harmful to sensitive plants. Do not mix with other chemicals, other than Terraclor®, unless prior use has proven compatibility.

RECOMMENDATIONS

For bedding and bench plantings use with water, as a soil drench at time of seeding and transplanting. The drench solution should be continually agitated to assure uniform application of the fungicidal material. To improve penetration of the fungicide, the drenching should be followed with an equal quantity of water, applied either by sprinkling or by irrigation.

Bedding Plants: Alyssum, Calendula, Chrysanthemum, Celosia, Dahlia, Dianthus, Marigold, Pansies, Petunia, Salvia, Snapdragon, Vinca, Verbena, Zinnia.

Foliage Plants: Aglaonema, Anthurium, Chinese Evergreen, Dieffenbachia, Nephthytis, Peperomia, Pilea, Pothos, Philodendron Spp.

Herbaceous and Woody: African Violet, Geranium, Rhododendron

BEDDING PLANTS: (Soil 2" to 3" deep):- Add 3/4 to 1 oz. to 25 gallons of water and apply to 100 sq. ft. of bed area. Repeat application may be made at one to two month intervals, if necessary.

TROPICAL FOLIAGE, HERBACEOUS and WOODY PLANTS: (Peat and other high organic media, 6" deep)-
Bed and Bench Plantings: Add 1 to 2 ozs. in 25 gallons of water and apply to 100 sq. ft. of bed or bench area. Repeat application at two to three month intervals to maintain disease control.

Pot Plantings: Use 1/2 pint solution per 6" pot, or in sufficient amount to saturate the soil.

USE CAUTIONS

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DO NOT PLANT FOOD CROPS IN ROTATION ON TREATED SOILS.

DESTROY THIS CONTAINER WHEN EMPTY

Terrazole®, Terraclor®, and Olin are trademarks of Olin Corporation

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.