

SODIUM OMADINE[®] 5% AQUEOUS SOLUTION

INDUSTRIAL MICROBIOSTAT

ACTIVE INGREDIENT: Sodium 2-pyridinethiol 1 Oxide* 5%
INERT INGREDIENTS: 95%
*Sodium Pyrithione

WARNING

KEEP OUT OF REACH OF CHILDREN


HARMFUL IF SWALLOWED. AVOID CONTACT WITH SKIN AND EYES.
IN CASE OF SKIN OR EYE CONTACT, FLUSH WITH PLENTY OF WATER.
GET MEDICAL ATTENTION IF IRRITATION PERSISTS.

DIRECTIONS FOR AQUEOUS METAL COOLANT AND CUTTING FLUID SOLUTION:

To inhibit bacteria growth add an initial dose of approx. 115 ppm Sodium Omadine 5% (9.20 lbs. Sodium Omadine 5% to 10,000 lbs. of solution) to the solution and a subsequent maintenance dose of approx. 55 ppm (4.4 lbs. of Sodium Omadine 5% to 10,000 lbs. of solution) every 7 to 10 days or as needed.

NOTE: Rinse empty container thoroughly with water and discard it.

55 GALS. NET



EPA REG. No. 1258-972-860

EPA EST. 1258-NY-3

OLIN CORPORATION
OLIN CHEMICALS
DESIGNED PRODUCTS DEPARTMENT
120 LONG RIDGE ROAD, STAMFORD, CONN. 06904

SO 55 AS 374