



**pool chlorinating
Type E cartridge for feeders**

Active ingredient:

Trichloro-s-Triazinetrione	99%
Inert Ingredients	1%
Available Chlorine	89%

NET WT. 4 LBS. (1.8 kg.)

**KEEP OUT OF REACH OF CHILDREN
DANGER!**
See Usage Instructions and Precautions on Back Panel

EPA Reg. No. 1258-997 EPA Est. 9157-MI-1
For use in E-Z Clor, CPC and SUN Feeders

Olin CHEMICALS
WATER PRODUCTS AND SERVICES DIVISION
OLIN CORPORATION 120 Long Ridge Road, Stamford, CT. 06904

ACCEPTED
with COMMENTS
in EPA Letter Dated:

SEP 05 1986

Under the Federal Insecticide,
Fungicide, and Herbicide Act
as amended, for the pesticide
registered under EPA Reg. No.
1258-997

BEST AVAILABLE COPY

PACE

Pool Chlorinating Type E Cartridge for use in E-Z Chlor, C-C and C-4 Feeders

- Protects against bacteria and algae
- Stabilized (Sun Protector™) for extended chlorine life
- Slow dissolving
- Totally soluble
- The PACE Tablets in this cartridge will provide an effective chlorine residual of 1.0-1.5 parts per million (ppm) in 10,000 gallons of water for about 9 weeks under normal conditions.

INSTRUCTIONS FOR POOL USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use with E-Z Chlor, CPC and Sun Feeders.

START UP

1. Bring chlorine residual to 1.0-1.5 ppm. Follow instructions for your specific feeding device shown below.
2. The day after your chlorine residual reaches 1.0-1.5 ppm, use cyanuric acid (e.g., PACE™ Stabilizer - Concentrate) to stabilize your pool. Follow label directions.
3. Adjust pH to the ideal range of 7.2-7.6.

INSTRUCTIONS FOR USE IN E-Z CHLOR AND SUN FEEDERS

1. Remove and save these instructions for future reference.
2. Make sure that the cartridge lid is tightly secured.
3. Punch out all of the dimples on the bottom of the cartridge (4).
4. Punch out 1 dimple on the bottom of each side (in curved area) (2).
5. Use the following table and punch out 1 hole on 1 flat side as noted.

Approx. Pool Size (sq-ft)

- 5,000 - lowest dimple
- 15,000 - second from bottom
- 25,000 - third from bottom
- 35,000 - fourth from bottom

6. If using two cartridges at a time, divide the pool size by 2 to determine what holes should be punched in the cartridge (e.g. for 2 cartridges in 30,000 gallons, use the directions searched for 15,000 gallons).
 7. Place the cartridge in the feeder.
 8. Maintain the chlorine residual between 1.0 and 1.5 ppm. Test the water frequently using a suitable test kit.
- If the chlorine level is below 1.0, punch out a higher dimple. If the chlorine residual is above 1.5, punch out a lower dimple and cover the original hole with black electrical tape.
- Periodically remove the cartridge and shake it to determine the approximate tablet level. Insert a new cartridge when old one is empty.

INSTRUCTIONS FOR USE IN CPC FEEDER

1. Remove and save instructions for future use.
 2. Punch a 1/4" hole anywhere near the top of the cartridge.
 3. Punch 1/4" holes in the 4 dimples on the bottom of the cartridge.
 4. Punch 1/4" holes in the 4 dimples on the bottom curved sides of the cartridge.
 5. Place the cartridge in the feeder.
 6. Raise the calibration tube until the water level reaches the level corresponding to the number of gallons in your pool, per the feeder's instructions.
 7. Maintain the chlorine residual between 1.0 and 1.5 ppm. Test the quantity using a suitable test kit.
- If the chlorine residual is below 1.0, raise the calibration tube. If the chlorine residual is above 1.5, lower the calibration tube.
- Periodically remove the cartridge and shake it to determine the approximate tablet level. Insert a new cartridge when the old one is empty.

Precautionary Statements

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Highly corrosive. Causes skin and eye damage. May be fatal if swallowed. Do not get in eyes, on skin or on clothing. Do not handle with bare hands. Wear goggles or face shield and use rubber gloves or only thoroughly clean, dry utensils when handling. Irritating to nose and throat. Avoid breathing dust and fumes. Remove and wash contaminated clothing before reuse.

FIRST AID (Practical Treatment)

- If Swallowed:** Food large amounts of milk and/or water and induce vomiting. Call a physician immediately.
- If On Skin:** Brush off excess chemical and flush skin with cold water for at least 15 minutes. If irritation persists, get medical attention.
- If In Eyes:** Flush with cold water for at least 15 minutes. Call a physician immediately.
- If Inhaled:** Remove person to fresh air. Call a physician immediately.

CHEMICAL HAZARDS:

DANGER: Strong oxidizing agent. Use only clean, dry utensils. Mix only with water. Combustion with moisture, dirt, organic matter or other chemicals including other pool chemicals or any other foreign matter may start a chemical reaction with generation of heat, liberation of hazardous gases, and possible generation of fire and explosion. Avoid any contact with flammable or burning material such as a lighted cigarette. Do not use this cartridge in any chlorinated ing. device which has been in contact with any organic or prohibited chlorinating compounds (e.g. calcium hypochlorite). Such use may cause fire or explosion. Do not permit this cartridge to be in contact with plastic, contact plastic pool fittings or metal objects. Do not throw this cartridge directly into the pool.

ENVIRONMENTAL HAZARDS:

This product in the cartridge is toxic to fish. Do not discharge into lakes, streams, ponds, or public waters unless in accordance with an NPDES Permit. For guidance contact the regional office of EPA.

STORAGE AND DISPOSAL:

Keep product dry in original container. Store in a cool, dry, well ventilated area away from heat or open flame. Triple rinse empty cartridge thoroughly by immersing in pool water. Drain and then securely wrap original cartridge in several layers of newspaper and discard in trash.

EMERGENCY HANDLING: In case of contamination or decomposition do not reuse cartridge. If possible, seal cartridge in open, well ventilated area. Flood with large volumes of water. Dispose of contaminated material in an approved landfill area or bury in a safe place.

ADDITIONAL POOL CARE RECOMMENDATIONS:

pH - Maintaining pH in the ideal range of 7.2-7.6 is extremely important to the efficacy of your chlorine and swimming comfort. Test the pH often with a suitable test kit. Adjust pH as necessary, using products such as PACE pH Increaser™ or PACE pH Plus™.

Supercalcination - Supercalcination with a calcium hypochlorite product such as PACE Super Chlorinator & Shock Treatment™ is recommended to burn out the organic material your filter can't handle and keep your water sparkling clear. Use as directed on the label, especially during excessive heat, when bathing loads are high, when water becomes dull looking or hazy, when there is a "chlorine" like smell or excessive eye irritation, or after heavy rains.

Shock Treatment - If algae appear, shock treat your pool with a calcium hypochlorite product such as PACE Super Chlorinator & Shock Treatment™. Follow label directions. An alternate method is to add 25 PACE Tablets (approx. 14 oz.) in the skimmer for each 10,000 gallons of water. Run pump and filter until totally dissolved (approx. 1 1/2 days). Thoroughly clean pool by scrubbing surface of algae growth, vacuum and cycle through filter. Pool should not be used until chlorine residual is between 1.0-1.5 ppm.

TO DETERMINE POOL CAPACITY:

Rectangular Pool: Multiply length x width x average depth x 7.5 = pool capacity (U.S. gallons)
Round Pools: Multiply radius (1/2 diameter) x radius x average depth x 2.25 = pool capacity (U.S. gallons)

PACE 628 4 000

BEST AVAILABLE COPY