



P.O. Box 991
Little Rock, Arkansas

Terra-Flo 21

Flowable
SEED TREATMENT FUNGICIDE
(With Dye)

ACTIVE INGREDIENTS	
Pentachloronitrobenzene	200
5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole*	100
INERT INGREDIENTS	700
TOTAL	1000

*This Product is supervised and tested in cotton production. Contains 100% of the active ingredients and does not contain Ethoxychlor, DDT, or other toxic materials.

Terra-Flo is a trademark of Olin Corporation

EPA Reg. No. 1258-1001
EPA Est. 1,758 MS 1

NET
CONTENTS

55
GALLONS

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
KEEP OUT OF REACH OF CHILDREN

DANGER

Corrosive. Causes skin damage and eye irritation. May cause skin sensitization. Do not get in eyes, on skin or on clothing. Wear rubber gloves when handling. Harmful if swallowed, inhaled or absorbed through skin. Wash thoroughly all contaminated clothing before reuse. Do not contaminate food or feedstuffs.

STATEMENT OF PRACTICAL TREATMENT
(First Aid Procedures)

In case of skin or eye contact, flush affected area with water for 15 minutes. Get immediate medical attention. Contains petroleum hydrocarbons. If swallowed, do not induce vomiting. Get prompt medical attention.

ENVIRONMENTAL HAZARDS

This material is toxic to fish. Keep out of lakes, ponds and streams. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Controls Rhizoctonia, Pythium and Fusarium seed and soil-borne seedling diseases of cotton. Using suitable liquid or slurry seed treating equipment such as Panogen or Mist-O-Matic, apply Terra-Flo 21 at rates shown below. Seed should be sound, well cured and cleaned prior to treatment. Use only the recommended rates of product per 100 pounds of cottonseed.

Acid delinted 13 1/2 fl ounces (400cc)
Mechanically delinted 18 fl ounces (533cc)

At the recommended rates, apply Terra-Flo 21 undiluted directly from the drum to the seed. If desired, add one part water to five parts Terra-Flo 21 and recalibrate equipment so that recommended rates will still be applied. Slight agitation may be required to mix the material.

This treatment on cottonseed will give only temporary protection to the planted seeds and seedlings. A subsequent soil application of Terraclor Super-X is necessary for the control of later stages of pre-emergence and post-emergence damping-off.

Keep container tightly closed when not in use. Do not feed treated seeds to animals.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food, or feed by storage or disposal.

Container Disposal: Rinse containers and dispose of in an incinerator or landfill approved for pesticide containers.

General: Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.

NOTICE TO BUYER

The product conforms to its chemical description and is fit for the use intended when used in accordance with label directions. Olin will not be responsible for special, consequential or contingent damages arising from injury to persons or property.

LR1001-081

1988
12/15/88