

GENERAL INFORMATION

If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using.

When filling spray tank first fill tank 1/2 of water; add the calculated amount of 2,4-D emulsifiable, then finish filling tank with water with by-pass open and pump running.

Material may tend to settle when not agitated. When mixture is left in sprayer allow by-pass pressure to re-agitate before resuming spray.

PRECAUTION

BECAUSE OF THE DIFFICULTIES OF REMOVING 2,4-D and 2,4,5-T FROM SPRAYERS, DO NOT USE THE SAME SPRAY EQUIPMENT FOR APPLYING OTHER MATERIALS TO PLANTS AS INJURY WILL RESULT. USE A SEPARATE SPRAYER FOR THIS PURPOSE.

Do not reuse empty container. Destroy by burying in a safe place. Do not burn.

ACCEPTED

AMINE 40
WEED KILLER

ACTIVE INGREDIENT:

1258-114 Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid 49.5%
(Equivalent to 2,4-Dichlorophenoxyacetic Acid 41.1%)

INERT INGREDIENTS:

Total 50.5%
100.0%

Contains the equivalent of 4 lbs. of 2,4-Dichlorophenoxyacetic Acid per gallon at 68° F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION. Avoid contact with eyes, skin and clothing. Do not take internally. Do not contaminate feed or foodstuffs.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Course sprays are less likely to drift.)

Do not contaminate domestic or irrigating water supplies.

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not seed lawns, gardens or fields until at least four weeks after application of spray to area. Do not mix with liquid fertilizers. Do not apply in any manner not specified on the label.

DO NOT STORE NEAR FERTILIZERS, SEEDS, INSECTICIDES, FUNGICIDES OR PLANTS

CROP	*Amount Amine 40 Per Acre	Equivalent 2,4-D Per Acre	APPLICATION	
CORN, MILO	1 pt.	1/2 lb.	When crop is small and more resistant to 2,4-D but when weeds are succulent or in active growth stage.	Use drop nozzles. Do not spray after
WHEAT	1 - 1 1/2 pt.	1/2 - 3/4 lb.	FALL In spring when weeds are small and grain is in full tiller stage (4 to 8 inches high) and always before boot stage.	Do not apply during grain fields, within 100 yds. to livestock.
OATS			SPRING After plants are fully tillered but before stems begin to joint.	
BARLEY	1 - 1 1/3 pts.	1/2 - 3/4 lb.		Some varieties of Sheyenne, Royal, Ko
RYE			When plants have formed 4 or 5 leaves but before plants have come into bud or bloom stage.	Injury may result to from early spring stage. Do not apply graze treated areas
FLAX	1/2 pt.	1/4 lb.		
PASTURES, MEADOWS	1 - 2 qts.	2 - 4 lbs.	On established stands of perennial grasses apply in early spring when sufficient weeds have emerged and are actively growing and not too mature.	Spray 3 to 5 days after mowing. Retreatment may cause injury to bent grass, hair grass, ornamentals from spray drift.
LAWNS, ORNAMENTAL TURF GRASS, GOLF FAIRWAYS	Large areas - 2 pints per acre. Small areas - 2 tablespoonfuls in 3 to 5 gallons of water applied to 1000 square feet.			

*Application rates may vary slightly from State to State depending upon soil types and weather factors. Consult your State agricultural experiment station for recommendations under local conditions.

