

AMINE 40

WEED KILLER

ACTIVE INGREDIENT:
 Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid 49.5%
 (Equivalent to 2,4-Dichlorophenoxyacetic Acid 41.1%)

INERT INGREDIENTS: 50.5%
Total 100.0%

Contains the equivalent of 4 lbs. of 2,4-Dichlorophenoxyacetic Acid per gallon at 68° F.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION. Avoid contact with eyes, skin and clothing. Do not take internally. Do not contaminate feed or foodstuffs.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Course sprays are less likely to drift.)

Do not contaminate domestic or irrigating water supplies.

Vapors from this product may injure susceptible plants in the immediate vicinity.

Do not seed lawns, gardens or fields until at least four weeks after application of spray to area. Do not mix with liquid fertilizers. Do not apply in any manner not specified on the label.

DO NOT STORE NEAR FERTILIZERS, SEEDS, INSECTICIDES, FUNGICIDES OR PLANTS

TIME OF APPLICATION

Annual weeds are most susceptible in the seedling stage and early growth stages of development and when conditions are such as to promote a vigorous growth. WHEN USED IN CROP SPRAYING AS A SELECTIVE SPRAY, THE STAGE OF GROWTH OF THE CROP MUST ALSO BE CONSIDERED.

SOME SUSCEPTIBLE WEEDS

- | | | |
|--------------|----------------|-----------------------|
| Arrowhead | Henbit | Prickly Lettuce |
| Artichoke | Horsetail | Puncture Vine |
| Beggar Ticks | Jimsonweed | Purslane |
| Ball Thistle | Kochia | Ragweed |
| Butterprint | Lambs Quarters | Shepherd's Purse |
| Catnip | Loco Weed | Smart Weed |
| Chickory | Mallow | Sow Thistle, (annual) |
| Cocklebur | Marestail | Slinkweed |
| Dandelion | Mexican Weed | Sumacs |
| Docks | Morning Glory | Sunflower |
| Fanweed | Mustards | Wild Mustard |
| Figwort | Nettles | Wild Onion |
| Four O'Clock | Pepper Grass | Wild Parsnip |
| Ground Ivy | Pig Weed | Wild Radish |
| | Wild Rape | Wild Sweet Potato |

Some weeds above, listed as susceptible may not be completely killed but growth is so retarded by proper application that they are no longer a factor affecting growth and yield of the crop.

NOTICE TO BUYER

Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damage arising from a breach of this warranty.

ON ACCEPTED

APR 23 1969
 1258-1006
 1258-164

it may agitate
 1/4 16H
 f 2,4-D
 h water
 agitated.
 by-pass spray
 MOVING DO NOT FOR AP. ANTS AS SEPARATE
 destroy by

Equivalent 2,4-D Per Acre	APPLICATION	CAUTION
1/2 lb.	When crop is small and more resistant to 2,4-D but when weeds are succulent or in active growth stage.	Use drop nozzles when crop is 10 in. tall or higher. Keep spray off leaves. Do not spray after tassels appear. Some injury to corn may occur.
FALL pts. 1/2 - 3/4 lb.	In spring when weeds are small and grain is in full tiller stage (4 to 8 inches high) and always before boot stage.	Do not apply during boot or dough stage. Do not forage or graze treated grain fields within two weeks after treatment. Do not feed treated straw to livestock.
SPRING pts. 1/2 - 3/4 lb.	After plants are fully tilled but before stems begin to joint.	Some varieties should not be sprayed (tolerant varieties: Redwing, Sheyenne, Royal, Kola, Dakota). Do not overdose.
1/4 lb.	When plants have formed 4 or 5 leaves but before plants have come into bud or bloom stage.	Injury may result to bent grass and to clovers. Usually clovers will recover from early spring applications. Do not apply when grass is in boot to milk stage. Do not apply to newly seeded areas or after heading begins. Do not graze treated areas to dairy animals within 7 days after treatment.
1/2 - 4 lbs.	On established stands of perennial grasses apply in early spring when sufficient weeds have emerged and are actively growing and not too mature.	

Spray 3 to 5 days after mowing. Retreatment of perennial weed areas may be necessary. May cause injury to bent grass, carpet grass, dichondra, St. Augustine and to clovers. Protect ornamentals from spray drift.

For areas — 2 pints per acre.
 For all areas — 2 tablespoonfuls in 3 to 5 gallons of water applied to 1000 square feet.

Consult your State agricultural experiment station for recommendations under local conditions or more specific uses.

LR 87K-035 6224334 USDA Reg. No. 1258-164

