



100% CONTROL
OF ROOT ROT

ACCEPTED

JUN 10 1982

U.S. Federal Insecticide,
Fungicide, and Rodenticide Act,
approved for the pesticide
registration number

EPA Reg. No. 1258-1029

Terrazole

5% GRANULAR
FUNGICIDE

ACTIVE INGREDIENT:
5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole* 5.0%
INERT INGREDIENTS: 95.0%
TOTAL 100.0%

*U.S. Patents 3,260,568 and 3,260,725, others pending

Terrazole is a trademark of Olin Corporation

EPA Reg. No. 1258-1029
EPA Est. 125R-MS-1

PRECAUTIONARY STATEMENTS **Hazards to Humans and Domestic Animals** **KEEP OUT OF REACH OF CHILDREN**

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not handle or apply product with bare hands.

STATEMENT OF PRACTICAL TREATMENT **(First Aid Procedures)**

If swallowed, induce vomiting. If on skin, remove contaminated clothing and wash skin with soap and warm water. If inhaled, remove to fresh air. If in eyes, flush with water for 15 minutes.

For further treatment, get medical attention.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes.
Do not apply directly to lakes, streams or ponds.

STORAGE AND DISPOSAL

PROHIBITIONS Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL Pesticide that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL **PAPER BAGS** - Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act. **PLASTIC CONTAINERS** - Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if permitted by State and local authorities.

GENERAL Consult federal, state or local disposal authorities for approved method of disposal. Do not use as limited open burning.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

ORNAMENTALS - For control of damping-off and root rot diseases caused by Pythium and Phytophthora at the time of seeding or transplanting.

PLANT TYPE		RECOMMENDATIONS
FOLIAGE PLANTS		Dry Soil Mix
Aglaonema Anthurium Brassia Caladium Chinese Evergreen Daffodorus Dracena	Echeveria Epiphyllum Nepenthes Peperomia Pilea Philodendron Zygocactus	To mix with potting, greenhouse or other transplanting soils, add Terrazole 5G at the rate of 10 ozs. per cubic yard of soil and mix thoroughly. Soil to be mixed should be free of large lumps and dry enough to prevent agglomeration during mixing. Where Rhizogonium control is desired, add 100 lbs. of 0.32% Terrazole 5G to 100 lbs. of soil. Retreat as needed for continued control with soil drench applications of Terrazole 35% Wettable Powder or Terrazole 25% Emulsifiable.
CONTAINER & BEDGROWN		
HERBACEOUS	WOODY	NOTE: Overdosage may result in phytotoxicity. If plants are suspected of being sensitive, treat only a few plants and observe for 4 to 5 days prior to treating the entire planting.
African Violet Begonia Calceolaria Chrysanthemum Easter Lily Geranium Gerbera Gladiolus Gypsophila Orchids Sinningia	Arctostaphylos Azalea Camelia Ceanothus Ficus Fir Hemlock Holly Juniper Leucothoe Magnolia Poinsettia Rhamnus Rhododendron Southern Pine Western Pine	

AVOCADOS (California Only) For control of Avocado root rot (Phytophthora cinnamomi). Apply Terrazole 5G Granular evenly to the soil surface surrounding the Avocado tree, in a circular area extending out to the drip line. The amount of material to be applied to each tree will be governed by the area to be covered, as per chart below.

Diameter of Circular Surface	Approx. Sq. Ft. Covered	Amount of Terrazole 5G Gran. per tree
20' dia.	314	86 ounces
15' dia.	176	48 ounces
10' dia.	79	22 ounces
5' dia.	20	6 ounces

Following application apply approx. one gallon of water per square foot of area treated. Repeat application once a month for 9 to 10 months during the growing season.

Do not treat within 30 days of harvest. During application, care should be exercised so that the treatment does not come in contact with the fruit.

LR1029-05F

NET
CONTENTS