

1/12/93

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warning and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

*DACONIL is a registered trademark of ISK Biotech.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

CHIPCO is a registered trademark of Rhône-Poulenc.
©1991 Rhône-Poulenc Ag Company

CHIPCO® brand 26019 Fungicide (PENDING) Submitted 1/11/93.

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

AUG 5 1993

Ms. Judith K. Lemm
 Rockland Corporation
 P. O. Box 809
 West Caldwell, NJ 07006

Dear Ms. Lemm:

Subject: Revised Labeling
 Benomyl Fungicide
 EPA Registration No. 572-254
 Your Application Dated July 22, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable. A stamped accepted copy for the label is enclosed for your record.

Please submit five(5) copies of your final printed labeling.

If you have any questions regarding this matter, please contact me or Sidney Jackson at (703) 305-7610.

Sincerely yours,

col

Clarence O Lewis, III
 Acting Product Manager (21)
 Fungicide-Herbicide Branch
 Registration Division (H7505C)

Enclosure

CONCURRENCES

SYMBOL	H7505C					
SURNAME	JACKSON	C. LEWIS				
DATE	8/5/93	8/5/93				

2 9 3

Rockland. Benomyl Fungicide

- A systemic fungicide to be mixed with water for spray application
- Controls listed diseases on Turf, and certain fruits and vegetables
- Provides both curative (eradicant) and protective action

ACTIVE INGREDIENT:

Benomyl (Methyl 1-(butylcarbamoyl)
2-benzimidazolecarbamate)50%
INERT INGREDIENTS:50%
100%

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside for additional precautionary statements
BE SURE AND READ THE COMPLETE PACKAGE LABEL
AND THIS LEAFLET BEFORE USING THIS PRODUCT.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Get medical attention.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Avoid alcohol.

IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water. Launder clothing before reuse.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION
May be irritating to eyes, nose, throat and skin. May cause temporary allergic skin reactions in a few susceptible persons which should be treated as allergic dermatitis. There is no evidence of after effects or permanent injury. Avoid contact with skin, eyes and clothing. Avoid breathing dust or spray. When using, wear chemical resistant gloves, long pants and long sleeved shirt. Apply with wind to your back. Wash non-disposable gloves with soap and water before removing. Do not enter or allow reentry into treated areas until sprays have dried unless wearing the protective clothing specified.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish in adjacent areas. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARD: Keep away from fire or sparks.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply as a spray using sufficient water to obtain thorough coverage of the plants. Under severe disease conditions (when entire leaf is infected) use the higher rate and shorter interval specified for each application. Addition of ROCKLAND SPREADER STICKER to spray mixture enhances curative action and improves distribution of spray on hard-to-wet plants. Consult your State agricultural extension specialist for guidance on your particular disease control situation.

TURF: Apply 6 Tablespoons in 1-2 1/2 gallons of water per 500 sq. ft. for Dollar Spot, Large Brown Patch. Apply at earliest appearance of disease and continue at 10-14 day intervals as long as needed. Shorten interval to 5-7 days when conditions are unusually



BEST AVAILABLE COPY

3 9 3

favorable for disease development. For Strip Smut use 18 Tablespoons in 1-2 1/2 gallons of water for 500 sq. ft. in October or early Spring before grass begins to grow. After application, water turf sufficiently to carry product to base of plants.

TOMATOES (Garden): Apply 1/2-1 Tbsp. per gallon of water to control Gray (Botrytis), Leaf (Cladosporium) and White (Sclerotinia) Molds, Cercospora Leaf Spot, Phoma Leaf Spot. Begin applications when disease first appears and repeat at 7-14 day intervals as needed.

BEANS (Snap and Lima): Use 1 1/2-2 Tablespoons per gallon of water for White (Sclerotinia) and Gray (Botrytis) Molds. Begin at 1/4 to 1/2 bloom; repeat at peak bloom. Do not apply to Snap or Dry Beans within 14 days of harvest, nor to Lima Beans within 28 days of harvest.

CUCUMBERS, MELONS, PUMPKINS, SQUASH:

Use 1/4 to 1/2 Tablespoon per gallon of water for Target Spot (Cucumbers), Gummy Stem Blight, Powdery Mildew, Anthracnose. Begin applications when plants begin to run or when disease first appears; repeat at 7-14 day intervals as needed. For Target Spot, use 7 day intervals as needed.

FRUIT TREES: For Pears, Apples, use 1/4-1/3 Tablespoon per gallon of water for Powdery Mildew, Scab, Sooty Blotch, Flyspeck. Apply at 1/2 green tip and repeat at 7-14 day intervals (or as needed) through cover sprays. For Peaches, Cherries, treatment is most effective if applied just before rainfall. Use 1/2 Tablespoon per gallon of water. For Brown Rot, Blossom Blight, apply on Peaches at pink bud; on Cherries at early popcorn. Make second application at 3/4-full bloom. If blossoming is prolonged or favorable disease conditions persist, apply at petal fall. For Fruit Brown Rot, after Blossom Blight sprays make 2 preharvest applications beginning 3 weeks before harvest up to day of harvest. For Peach Scab, Powdery Mildew, use Blossom Blight spray schedule plus applications at shuck, split, shuck fall and 14 days later. For Cherry Leaf Spot, use Blossom Blight spray schedule and continue at 10-14 day intervals through harvest. Make an additional application 2-3 weeks after harvest. To help control resistant strains of fungi on Apples, Peaches, add 3/4 tsp. of ROCKLAND CAPTAN FUNGICIDE per gallon of spray; on Cherries, add 3/4 tsp. of ROCKLAND CAPTAN FUNGICIDE per gallon of spray

This package is sold by weight, not by volume. Some settling of contents may have occurred during shipping and handling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store unused material, in original container only, in a locked storage area out of reach of children and animals. Keep container tightly closed. Never allow product to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness as a fungicide.

DISPOSAL: Do not reuse empty container. Securely wrap in several layers of newspaper and discard in trash.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 572-254
Distributed by
Rockland Corporation
West Caldwell, NJ 07006

EPA Est. No. 572-NJ-1

1091

BEST AVAILABLE COPY